

## **RING1** Antibody

Rabbit mAb Catalog # AP92237

## **Product Information**

Application	WB, IHC, IF, ICC, IHF
Primary Accession	<u>Q06587</u>
Reactivity	Human
Clonality	Monoclonal
Other Names	Ring1A; Ring1; RING1A; Rnf1;
lsotype	Rabbit IgG
Host	Rabbit
Calculated MW	42429

## **Additional Information**

Dilution	WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human RING1
Description	Constitutes one of the E3 ubiquitin-protein ligases that mediate
	monoubiquitination of 'Lys-119' of histone H2A, thereby playing a central role
	in histone code and gene regulation. H2A 'Lys-119' ubiquitination gives a
	specific tag for epigenetic transcriptional repression and participates in X
	chromosome inactivation of female mammals.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium
	azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term.
	Avoid freeze / thaw cycle.

## **Protein Information**

Name	RING1 ( <u>HGNC:10018</u> )
Function	Constitutes one of the E3 ubiquitin-protein ligases that mediate monoubiquitination of 'Lys-119' of histone H2A, thereby playing a central role in histone code and gene regulation. H2A 'Lys-119' ubiquitination gives a specific tag for epigenetic transcriptional repression and participates in X chromosome inactivation of female mammals. Essential component of a Polycomb group (PcG) multiprotein PRC1-like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development. PcG PRC1 complex acts via chromatin remodeling and modification of histones, rendering chromatin heritably changed in its expressibility. Compared to RNF2/RING2, it does not have the main E3 ubiquitin ligase activity on histone H2A, and it may rather act as a modulator of RNF2/RING2 activity.
Cellular Location	Nucleus. Nucleus speckle



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.