

MCAK Antibody

Rabbit mAb

Catalog # AP92273

Product Information

Application	WB, IF, FC, ICC
Primary Accession	Q99661
Reactivity	Human
Clonality	Monoclonal
Other Names	MCAK; KNSL6;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	81313

Additional Information

Dilution	WB 1:500~1:2000 ICC/IF 1:50~1:200 FC 1:100
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human MCAK
Description	Promotes ATP-dependent removal of tubulin dimers from microtubules. Regulates the turnover of microtubules at the kinetochore and functions in chromosome segregation during mitosis.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.


Protein Information

Name	KIF2C
Synonyms	KNSL6
Function	In complex with KIF18B, constitutes the major microtubule plus-end depolymerizing activity in mitotic cells (PubMed: 21820309). Regulates the turnover of microtubules at the kinetochore and functions in chromosome segregation during mitosis (PubMed: 19060894). Plays a role in chromosome congression and is required for the lateral to end- on conversion of the chromosome-microtubule attachment (PubMed: 23891108).
Cellular Location	Cytoplasm, cytoskeleton. Nucleus {ECO:0000250 UniProtKB:P70096} Chromosome, centromere. Chromosome, centromere, kinetochore. Note=Associates with the microtubule network at the growing distal tip (the plus-end) of microtubules, probably through interaction with MTUS2/TIP150 and MAPRE1 (By similarity). Association with microtubule plus ends is also mediated by interaction with KIF18B. Centromeric localization requires the presence of BUB1 and SGO2. {ECO:0000250 UniProtKB:P70096,

Tissue Location

Expressed at high levels in thymus and testis, at low levels in small intestine, the mucosal lining of colon, and placenta, and at very low levels in spleen and ovary; expression is not detected in prostate, peripheral blood Leukocytes, heart, brain, lung, liver, skeletal muscle, kidney or pancreas. Isoform 2 is testis- specific.

Images



Western blot analysis of MCAK expression in MCF-7 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.