

Legumain Antibody

Rabbit mAb

Catalog # AP92308

Product Information

Application	WB
Primary Accession	Q99538
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	AEP; cysteine 1; Legumain; LGMN; LGMN1; PRSC1;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	49411


Additional Information

Dilution	WB 1:500~1:2000
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Legumain
Description	Has a strict specificity for hydrolysis of asparaginyl bonds. Can also cleave aspartyl bonds slowly, especially under acidic conditions. May be involved in the processing of proteins for MHC class II antigen presentation in the lysosomal/endosomal system.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	LG MN {ECO:0000303 PubMed:30425301, ECO:0000312 HGNC:HGNC:9472}
Function	Has a strict specificity for hydrolysis of asparaginyl bonds (PubMed: 23776206). Can also cleave aspartyl bonds slowly, especially under acidic conditions (PubMed: 23776206). Involved in the processing of proteins for MHC class II antigen presentation in the lysosomal/endosomal system (PubMed: 9872320). Also involved in MHC class I antigen presentation in cross-presenting dendritic cells by mediating cleavage and maturation of Perforin-2 (MPEG1), thereby promoting antigen translocation in the cytosol (By similarity). Required for normal lysosomal protein degradation in renal proximal tubules (By similarity). Required for normal degradation of internalized EGFR (By similarity). Plays a role in the regulation of cell proliferation via its role in EGFR degradation (By similarity).
Cellular Location	Lysosome.
Tissue Location	Ubiquitous. Particularly abundant in kidney, heart and placenta.

Images



Western blot analysis of Legumain expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.