

GGT1 Antibody

Rabbit mAb Catalog # AP92408

Product Information

Application	WB, IHC
Primary Accession	<u>P19440</u>
Reactivity	Human
Clonality	Monoclonal
Other Names	GGT; GTG; CD224; GGT 1;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	61410

Additional Information

Dilution Purification	WB 1:500~1:2000 IHC 1:50~1:200 Affinity-chromatography
Immunogen	A synthesized peptide derived from human GGT1
Description	Initiates extracellular glutathione (GSH) breakdown, provides cells with a local cysteine supply and contributes to maintain intracelular GSH level. It is part of the cell antioxidant defense mechanism. Catalyzes the transfer of the glutamyl mojety of glutathione to amino acids and dipeptide acceptors.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	GGT1
Synonyms	GGT
Function	Cleaves the gamma-glutamyl bond of extracellular glutathione (gamma-Glu-Cys-Gly), glutathione conjugates (such as maresin conjugate (13R)-S-glutathionyl-(14S)-hydroxy-(4Z,7Z,9E,11E,16Z,19Z)- docosahexaenoate, MCTR1) and other gamma-glutamyl compounds (such as leukotriene C4, LTC4) (PubMed: <u>17924658</u> , PubMed: <u>21447318</u> , PubMed: <u>27791009</u>). The metabolism of glutathione by GGT1 releases free glutamate and the dipeptide cysteinyl-glycine, which is hydrolyzed to cysteine and glycine by dipeptidases (PubMed: <u>27791009</u>). In the presence of high concentrations of dipeptides and some amino acids, can also catalyze a transpeptidation reaction, transferring the gamma-glutamyl moiety to an acceptor amino acid to form a new gamma-glutamyl compound (PubMed: <u>17924658</u> , PubMed: <u>21447318</u> , PubMed: <u>7673200</u> , PubMed: <u>7759490</u> , PubMed: <u>8095045</u> , PubMed: <u>8827453</u>). Contributes to cysteine homeostasis, glutathione homeostasis and in the

	conversion of the leukotriene LTC4 to LTD4.
Cellular Location	Cell membrane; Single-pass type II membrane protein {ECO:0000250 UniProtKB:P07314}
Tissue Location	Detected in fetal and adult kidney and liver, adult pancreas, stomach, intestine, placenta and lung. There are several other tissue-specific forms that arise from alternative promoter usage but that produce the same protein

Images



Western blot analysis of GGT1 expression in Human fetal liver lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.