

PLCB3 Antibody

Rabbit mAb

Catalog # AP92439

Product Information

Application	WB, IHC
Primary Accession	Q01970
Reactivity	Human
Clonality	Monoclonal
Other Names	Phosphoinositide phospholipase C; Phospholipase C beta 3; PLC beta 3; Plcb3;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	138799


Additional Information

Dilution	WB 1:500~1:2000 IHC 1:50~1:200
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human PLCB3
Description	The production of the second messenger molecules diacylglycerol (DAG) and inositol 1,4,5-trisphosphate (IP3) is mediated by activated phosphatidylinositol-specific phospholipase C enzymes.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	PLCB3 {ECO:0000303 PubMed:20966218, ECO:0000312 EMBL:AAA77683.1}
Function	Catalyzes the production of the second messenger molecules diacylglycerol (DAG) and inositol 1,4,5-trisphosphate (IP3) (PubMed: 20966218 , PubMed: 29122926 , PubMed: 37991948 , PubMed: 9188725). Key transducer of G protein-coupled receptor signaling: activated by G(q)/G(11) G alpha proteins downstream of G protein-coupled receptors activation (PubMed: 20966218 , PubMed: 37991948). In neutrophils, participates in a phospholipase C-activating N-formyl peptide-activated GPCR (G protein-coupled receptor) signaling pathway by promoting RASGRP4 activation by DAG, to promote neutrophil functional responses (By similarity).
Cellular Location	Cytoplasm. Membrane {ECO:0000250 UniProtKB:Q99JE6}. Nucleus {ECO:0000250 UniProtKB:P51432} Note=And particulate fractions. {ECO:0000250 UniProtKB:Q99JE6}

Images



Western blot analysis of PLCB3 expression in A431 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.