

GRIK2 Antibody

Rabbit mAb

Catalog # AP92443

Product Information

Application	WB, IHC
Primary Accession	Q13002
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	EAA4; GLR6; MRT6; GLUK6; GLUR6; GluK2;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	102583

Additional Information

Dilution	WB 1:500~1:2000 IHC 1:50~1:200
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human GRIK2
Description	Ionotropic glutamate receptor. L-glutamate acts as an excitatory neurotransmitter at many synapses in the central nervous system. Binding of the excitatory neurotransmitter L-glutamate induces a conformation change, leading to the opening of the cation channel, and thereby converts the chemical signal to an electrical impulse.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.


Protein Information

Name	GRIK2
Synonyms	GLUR6
Function	<p>Ionotropic glutamate receptor that functions as a cation permeable ligand-gated ion channel, gated by L-glutamate and the glutamatergic agonist kainic acid. L-glutamate acts as an excitatory neurotransmitter at many synapses in the central nervous system. Binding of the excitatory neurotransmitter L-glutamate induces a conformation change, leading to the opening of the cation channel, and thereby converts the chemical signal to an electrical impulse. The receptor then desensitizes rapidly and enters a transient inactive state, characterized by the presence of bound agonist (PubMed:14511640, PubMed:28180184, PubMed:34375587, PubMed:7536611, PubMed:8730589). Modulates cell surface expression of NETO2. In association with GRIK3, involved in presynaptic facilitation of glutamate release at hippocampal mossy fiber synapses (By similarity).</p>

Cellular Location Cell membrane; Multi-pass membrane protein. Postsynaptic cell membrane {ECO:0000250|UniProtKB:P42260}; Multi-pass membrane protein

Tissue Location Expression is higher in cerebellum than in cerebral cortex.

Images



Western blot analysis of GRIK2 expression in A431 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.