

PLTP Antibody

Rabbit mAb

Catalog # AP92472

Product Information

Application	WB
Primary Accession	P55058
Reactivity	Rat, Human
Clonality	Monoclonal
Other Names	BPIFE; HDLCQ9; OD107; Pltp;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	54739

Additional Information

Dilution	WB 1:500~1:2000
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human PLTP
Description	Converts HDL into larger and smaller particles. May play a key role in extracellular phospholipid transport and modulation of hdl particles.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	PLTP
Function	Mediates the transfer of phospholipids and free cholesterol from triglyceride-rich lipoproteins (low density lipoproteins or LDL and very low density lipoproteins or VLDL) into high-density lipoproteins (HDL) as well as the exchange of phospholipids between triglyceride-rich lipoproteins themselves (PubMed: 11013307 , PubMed: 19321130 , PubMed: 21515415 , PubMed: 29883800 , PubMed: 7654777 , PubMed: 9132017). Facilitates the transfer of a spectrum of different lipid molecules, including diacylglycerol, phosphatidic acid, sphingomyelin, phosphatidylcholine, phosphatidylinositol, phosphatidylglycerol, cerebroside and phosphatidyl ethanolamine (PubMed: 9132017). Plays an important role in HDL remodeling which involves modulating the size and composition of HDL (PubMed: 29883800). Also plays a key role in the uptake of cholesterol from peripheral cells and tissues that is subsequently transported to the liver for degradation and excretion (PubMed: 21736953). Two distinct forms of PLTP exist in plasma: an active form that can transfer phosphatidylcholine from phospholipid vesicles to HDL, and an inactive form that lacks this capability (PubMed: 11013307).


Cellular Location

Secreted. Nucleus. Note=Nuclear export is XPO1/CRM1- dependent.

Tissue Location

Widely expressed. Highest level of expression in the ovary, thymus and placenta, with moderate levels found in the pancreas, small intestine, testis, lung and prostate. Low level expression in the kidney, liver and spleen, with very low levels found in the heart, colon, skeletal muscle, leukocytes and brain. Expressed in the cortical neurons.

Images



Western blot analysis of PLTP expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.