10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Oncostatin M Antibody

Rabbit mAb
Catalog # AP92514

Product Information

Application	WB
Primary Accession	P13725
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	Oncostatin M; OSM;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	28484

Additional Information

Dilution	WB 1:500~1:2000
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Oncostatin M
Description	Growth regulator. Inhibits the proliferation of a number of tumor cell lines. Stimulates proliferation of AIDS-KS cells. It regulates cytokine production, including IL-6, G-CSF and GM-CSF from endothelial cells.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.


Protein Information

Name	OSM (HGNC:8506)
Function	Functions as a cytokine that uses both type I OSM receptor (heterodimers composed of LIFR and IL6ST) and type II OSM receptor (heterodimers composed of OSMR and IL6ST) (PubMed:10997905, PubMed:39532904, PubMed:8999038). Functionally, regulates many processes including cell proliferation, cell differentiation, cytokine production (PubMed:10997905, PubMed:1542792, PubMed:1542793, PubMed:1717982, PubMed:2779549, PubMed:3540948). Mechanistically, low affinity ligand binding to IL6ST/gp130, induces heterodimerization with LIFR or OSMR, activating JAK tyrosine kinases (JAK1 or JAK2 and to a lesser extent TYK2) bound to their intracellular domains (PubMed:9188471). These kinases subsequently phosphorylate IL6ST/gp130 and LIFR or OSMR (PubMed:8999038). The tyrosine phosphorylated signaling receptors serve in turn as docking sites for recruitment and activation of STAT3 (PubMed:9188471). The type II OSM complex receptor is also able in addition to STAT3 to recruit STAT5B (PubMed:9188471). Moreover, the type II OSM complex receptor recruits SHC1 through OSMR in a JAK1 mediated phosphotyrosine-dependent manner, leading to SHC1 association with GRB2

and downstream activation of the Ras/Raf/MAPK pathway (PubMed:11016927).

Cellular Location Secreted

Images



Western blot analysis of Oncostatin M expression in Jurkat cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.