

DPYD Antibody

Rabbit mAb Catalog # AP92556

Product Information

Application WB, IHC, IF, FC, ICC, IHF

Primary Accession

Reactivity

Clonality

Q12882

Human

Monoclonal

Other Names DHP; DHPDHase; DPD; DPYD;

IsotypeRabbit IgGHostRabbitCalculated MW111401

Additional Information

Dilution WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200

Purification Affinity-chromatography

Immunogen A synthesized peptide derived from human DPYD

Description Involved in pyrimidine base degradation. Catalyzes the reduction of uracil and

thymine. Also involved the degradation of the chemotherapeutic drug

5-fluorouracil.

Storage Condition and Buffer Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium

azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term.

Avoid freeze / thaw cycle.

Protein Information

Name DPYD (HGNC:3012)

Function Involved in pyrimidine base degradation (PubMed: <u>1512248</u>). Catalyzes the

reduction of uracil and thymine (PubMed: 1512248). Also involved the

degradation of the chemotherapeutic drug 5-fluorouracil (PubMed: 1512248).


Cellular Location Cytoplasm.

Tissue Location Found in most tissues with greatest activity found in liver and peripheral

blood mononuclear cells

Images

Western blot analysis of DPYD expression in HeLa cell lysate.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.