

SLC27A4 / FATP4 Antibody

Rabbit mAb

Catalog # AP92592

Product Information

Application	WB, IF, FC, ICC
Primary Accession	Q6P1M0
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	ACSVL4; FATP4; Fatty acid transport protein 4; IPS; S27A4; Solute carrier family 27 member4;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	72064

Additional Information

Dilution	WB 1:500~1:2000 ICC/IF 1:50~1:200 FC 1:100
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human SLC27A4 / FATP4
Description	plays a role in the transport of long chain fatty acids across the plasma membrane. It has acyl-coA ligase activity for long chain and very long chain fatty acids.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	SLC27A4 (HGNC:10998)
Function	Mediates the levels of long-chain fatty acids (LCFA) in the cell by facilitating their transport across cell membranes (PubMed: 10518211 , PubMed: 12556534 , PubMed: 20448275 , PubMed: 21395585 , PubMed: 22022213). Appears to be the principal fatty acid transporter in small intestinal enterocytes (PubMed: 20448275). Also functions as an acyl-CoA ligase catalyzing the ATP-dependent formation of fatty acyl- CoA using LCFA and very-long-chain fatty acids (VLCFA) as substrates, which prevents fatty acid efflux from cells and might drive more fatty acid uptake (PubMed: 22022213 , PubMed: 24269233). Plays a role in the formation of the epidermal barrier. Required for fat absorption in early embryogenesis (By similarity). Probably involved in fatty acid transport across the blood barrier (PubMed: 21395585). Indirectly inhibits RPE65 via substrate competition and via production of VLCFA derivatives like lignoceroyl-CoA. Prevents light-induced degeneration of rods and cones (By similarity).


Cellular Location

Endoplasmic reticulum membrane; Multi-pass membrane protein

Tissue Location

Expressed at highest levels in brain, testis, colon and kidney. Expressed at medium levels in heart and liver, small intestine and stomach. Expressed at low levels in peripheral leukocytes, bone marrow, skeletal muscle and aorta. Expressed in adipose tissue (PubMed:24269233, PubMed:9878842). Expressed in brain gray matter (PubMed:21395585).

Images



Western blot analysis of SLC27A4 / FATP4 expression in (1) HeLa cell lysate; (2) Mouse kidney lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.