

TAF1 Antibody

Rabbit mAb

Catalog # AP92607

Product Information

Application	WB
Primary Accession	P21675
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	BA2R; CCG1; CCGS; DYT3; N TAF1; NSCL2; p250; TAF1; TAF2A; TAFII250; XDP;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	214714

Additional Information

Dilution	WB 1:500~1:2000
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human TAF1
Description	Largest component and core scaffold of the TFIID basal transcription factor complex. Contains novel N- and C-terminal Ser/Thr kinase domains which can autophosphorylate or transphosphorylate other transcription factors.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information


Name	TAF1 (HGNC:11535)
Synonyms	BA2R, CCG1, CCGS, TAF2A
Function	<p>The TFIID basal transcription factor complex plays a major role in the initiation of RNA polymerase II (Pol II)-dependent transcription (PubMed:33795473). TFIID recognizes and binds promoters with or without a TATA box via its subunit TBP, a TATA-box-binding protein, and promotes assembly of the pre-initiation complex (PIC) (PubMed:33795473). The TFIID complex consists of TBP and TBP-associated factors (TAFs), including TAF1, TAF2, TAF3, TAF4, TAF5, TAF6, TAF7, TAF8, TAF9, TAF10, TAF11, TAF12 and TAF13 (PubMed:33795473). TAF1 is the largest component and core scaffold of the TFIID complex, involved in nucleating complex assembly (PubMed:25412659, PubMed:27007846, PubMed:33795473). TAF1 forms a promoter DNA binding subcomplex of TFIID, together with TAF7 and TAF2 (PubMed:33795473). Contains novel N- and C-terminal Ser/Thr kinase domains which can autophosphorylate or transphosphorylate other transcription factors (PubMed:25412659, PubMed:8625415). Phosphorylates</p>

TP53 on 'Thr-55' which leads to MDM2- mediated degradation of TP53 (PubMed:[25412659](#)). Phosphorylates GTF2A1 and GTF2F1 on Ser residues (PubMed:[25412659](#)). Possesses DNA-binding activity (PubMed:[25412659](#)). Essential for progression of the G1 phase of the cell cycle (PubMed:[11278496](#), PubMed:[15053879](#), PubMed:[2038334](#), PubMed:[8450888](#), PubMed:[8625415](#), PubMed:[9660973](#), PubMed:[9858607](#)). Exhibits histone acetyltransferase activity towards histones H3 and H4 (PubMed:[15870300](#)).

Cellular Location

Nucleus

Images



Western blot analysis of TAF1 expression in SH-SY5Y cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.