

NUP62 Antibody

Rabbit mAb

Catalog # AP92630

Product Information

Application	WB
Primary Accession	P37198
Reactivity	Human
Clonality	Monoclonal
Other Names	IBSN; MGC841; NUP62; p62; SNDI;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	53255


Additional Information

Dilution	WB 1:500~1:2000
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human NUP62
Description	The nuclear pore complex is a structure that extends across the nuclear envelope and regulates the flow of macromolecules between the cytoplasm and the nucleus. Nucleoporins are the main components of the nuclear pore complex in eukaryotic cells.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	NUP62
Function	Essential component of the nuclear pore complex (PubMed: 1915414). The N-terminal is probably involved in nucleocytoplasmic transport (PubMed: 1915414). The C-terminal is involved in protein-protein interaction probably via coiled-coil formation, promotes its association with centrosomes and may function in anchorage of p62 to the pore complex (PubMed: 1915414 , PubMed: 24107630). Plays a role in mitotic cell cycle progression by regulating centrosome segregation, centriole maturation and spindle orientation (PubMed: 24107630). It might be involved in protein recruitment to the centrosome after nuclear breakdown (PubMed: 24107630).
Cellular Location	Nucleus, nuclear pore complex. Cytoplasm, cytoskeleton, spindle pole. Nucleus envelope. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Note=Central region of the nuclear pore, within the transporter (PubMed:1915414). During mitotic cell division, it associates with the poles of the mitotic spindle (PubMed:24107630)

Images



Western blot analysis of NUP62 expression in Raji cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.