

CD147 Antibody

Rabbit mAb Catalog # AP92645

Product Information

Application Primary Accession Reactivity Clonality Other Names	WB, IHC, IF, ICC, IHF <u>P35613</u> Human Monoclonal 5A11; 5F7; Basigin; Bsg; CD147; Collagenase stimulatory factor; EMMPRIN; M6 leukocyte activation antigen; Neurothelin; OK blood group antigen; TCSF; Tumor cell derived collagenase stimulatory factor;
lsotype	Rabbit IgG
Host	Rabbit
Calculated MW	42200

Additional Information

Dilution Purification	WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 Affinity-chromatography A synthesized peptide derived from CD147
Immunogen Description	Plays pivotal roles in spermatogenesis, embryo implantation, neural network
Storage Condition and Buffer	formation and tumor progression. Stimulates adjacent fibroblasts to produce matrix metalloproteinases (MMPS).

Protein Information

Name	BSG (<u>HGNC:1116</u>)
Function	[Isoform 1]: Essential for normal retinal maturation and development (By similarity). Acts as a retinal cell surface receptor for NXNL1 and plays an important role in NXNL1-mediated survival of retinal cone photoreceptors (PubMed:25957687). In association with glucose transporter SLC16A1/GLUT1 and NXNL1, promotes retinal cone survival by enhancing aerobic glycolysis and accelerating the entry of glucose into photoreceptors (PubMed:25957687). May act as a potent stimulator of IL6 secretion in multiple cell lines that include monocytes (PubMed:21620857).
Cellular Location	Melanosome. Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV. [Isoform 2]: Cell membrane; Single-pass type I membrane protein {ECO:0000250 UniProtKB:P26453}. Endosome Endoplasmic reticulum membrane; Single- pass type I membrane protein {ECO:0000250 UniProtKB:P26453} Basolateral cell membrane; Single-pass

	type I membrane protein {ECO:0000250 UniProtKB:P26453} [Isoform 4]: Cell membrane; Single-pass type I membrane protein {ECO:0000250 UniProtKB:P26453}
Tissue Location	[Isoform 1]: Retina-specific (PubMed:25957687). Expressed in retinal cone photoreceptors (at protein level) (PubMed:25957687). [Isoform 3]: Highly expressed in the bone marrow, fetal liver, lung, testis and thymus.

Images



Western blot analysis of CD147 expression in Jurkat cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.