

# Histone H3 (mono methyl K9) Antibody

Rabbit mAb Catalog # AP92647

### **Product Information**

**Application** WB, IHC, IF, ICC, IP, IHF

Primary Accession P68431

**Reactivity** Rat, Human, Mouse

**Clonality** Monoclonal

Other Names Histone H3.1, Histone H3, HIST1H3A;

IsotypeRabbit IgGHostRabbitCalculated MW15404

## **Additional Information**

**Dilution** WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 CHIP 1:50

**Purification** Affinity-chromatography

ImmunogenA synthesized peptide derived from Histone H3 (mono methyl K9)DescriptionBelongs to the histone H3 family. Play a central role in transcription

regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications

of histones, also called histone code, and nucleosome remodeling.

Storage Condition and Buffer Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium

azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term.

Avoid freeze / thaw cycle.

#### **Protein Information**

Name H3C1 ( <u>HGNC:4766</u>)

**Synonyms** H3FA, HIST1H3A


**Function** Core component of nucleosome. Nucleosomes wrap and compact DNA into

chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications

of histones, also called histone code, and nucleosome remodeling.

**Cellular Location** Nucleus. Chromosome.

# **Images**



Western blot analysis of Histone H3 (mono methyl K9) expression in Hela cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.