

Histone H3 (acetyl K18) Antibody

Rabbit mAb

Catalog # AP92648

Product Information

Application	WB, IHC, IF, FC, ICC, IP, IHF
Primary Accession	P68431
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	Histone H3.1, Histone H3, HIST1H3A;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	15404


Additional Information

Dilution	WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 CHIP 1:50 FC 1:50
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from Histone H3 (acetyl K18)
Description	Belongs to the histone H3 family. Play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	H3C1 (HGNC:4766)
Synonyms	H3FA, HIST1H3A
Function	Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.
Cellular Location	Nucleus. Chromosome.

Images



Western blot analysis of Histone H3 (acetyl K18)
expression in HeLa cell treated with TSA cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.