

Caspase-1 Antibody

Rabbit mAb

Catalog # AP92654

Product Information

Application	WB, IP
Primary Accession	P29466
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	CASP-1; ICE; IL-1 beta converting enzyme; IL-1BC; IL1BCE; caspase-1;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	45159

Additional Information

Dilution	WB 1:500~1:2000 IP 1:50
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from Caspase-1
Description	Thiol protease that cleaves IL-1 beta between an Asp and an Ala, releasing the mature cytokine which is involved in a variety of inflammatory processes. Important for defense against pathogens. Cleaves and activates sterol regulatory element binding proteins (SREBPs). Can also promote apoptosis.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	CASP1
Synonyms	IL1BC, IL1BCE
Function	Thiol protease involved in a variety of inflammatory processes by proteolytically cleaving other proteins, such as the precursors of the inflammatory cytokines interleukin-1 beta (IL1B) and interleukin 18 (IL18) as well as the pyroptosis inducer Gasdermin-D (GSDMD), into active mature peptides (PubMed: 15326478 , PubMed: 15498465 , PubMed: 1574116 , PubMed: 26375003 , PubMed: 32051255 , PubMed: 37993714 , PubMed: 7876192 , PubMed: 9334240). Plays a key role in cell immunity as an inflammatory response initiator: once activated through formation of an inflammasome complex, it initiates a pro-inflammatory response through the cleavage of the two inflammatory cytokines IL1B and IL18, releasing the mature cytokines which are involved in a variety of inflammatory processes (PubMed: 15326478 , PubMed: 15498465 , PubMed: 1574116 , PubMed: 32051255 , PubMed: 7876192). Cleaves a tetrapeptide after an Asp residue at position P1 (PubMed: 15498465 ,

PubMed:[1574116](#), PubMed:[7876192](#)). Also initiates pyroptosis, a programmed lytic cell death pathway, through cleavage of GSDMD (PubMed:[26375003](#)). In contrast to cleavage of interleukin IL1B, recognition and cleavage of GSDMD is not strictly dependent on the consensus cleavage site but depends on an exosite interface on CASP1 that recognizes and binds the Gasdermin-D, C-terminal (GSDMD-CT) part (PubMed:[32051255](#), PubMed:[32109412](#), PubMed:[32553275](#)). Cleaves and activates CASP7 in response to bacterial infection, promoting plasma membrane repair (PubMed:[22464733](#)). Upon inflammasome activation, during DNA virus infection but not RNA virus challenge, controls antiviral immunity through the cleavage of CGAS, rendering it inactive (PubMed:[28314590](#)). In apoptotic cells, cleaves SPHK2 which is released from cells and remains enzymatically active extracellularly (PubMed:[20197547](#)).


Cellular Location

Cytoplasm. Cell membrane

Tissue Location

Expressed in larger amounts in spleen and lung. Detected in liver, heart, small intestine, colon, thymus, prostate, skeletal muscle, peripheral blood leukocytes, kidney and testis. No expression in the brain.

Images



Western blot analysis of Caspase-1 expression in NIH/3T3 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.