

# UCHL3 Antibody

Rabbit mAb Catalog # AP92669

### **Product Information**

| Application       | WB, IHC                           |
|-------------------|-----------------------------------|
| Primary Accession | <u>P15374</u>                     |
| Reactivity        | Rat, Human, Mouse                 |
| Clonality         | Monoclonal                        |
| Other Names       | Ubiquitin thioesterase L3; UCHL3; |
| lsotype           | Rabbit IgG                        |
| Host              | Rabbit                            |
| Calculated MW     | 26183                             |


#### **Additional Information**

| Dilution<br>Purification<br>Immunogen    | WB 1:500~1:2000 IHC 1:50~1:200<br>Affinity-chromatography<br>A synthesized peptide derived from UCHL3  |
|--|--|
| Description Storage Condition and Buffer | Deubiquitinating enzyme (DUB) that controls levels of cellular ubiquitin<br>through processing of ubiquitin precursors and ubiquitinated proteins. Thiol<br>protease that recognizes and hydrolyzes a peptide bond at the C-terminal<br>glycine of either ubiquitin or NEDD8.<br>Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium<br>azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. |
|  | Avoid freeze / thaw cycle.   |

#### **Protein Information**

| Name              | UCHL3   |
|-------------------|---|
| Function          | Deubiquitinating enzyme (DUB) that controls levels of cellular ubiquitin<br>through processing of ubiquitin precursors and ubiquitinated proteins. Thiol<br>protease that recognizes and hydrolyzes a peptide bond at the C-terminal<br>glycine of either ubiquitin or NEDD8. Has a 10-fold preference for Arg and Lys<br>at position P3", and exhibits a preference towards 'Lys-48'-linked ubiquitin<br>chains. Deubiquitinates ENAC in apical compartments, thereby regulating<br>apical membrane recycling. Indirectly increases the phosphorylation of IGFIR,<br>AKT and FOXO1 and promotes insulin-signaling and insulin-induced<br>adipogenesis. Required for stress-response retinal, skeletal muscle and germ<br>cell maintenance. May be involved in working memory. Can hydrolyze<br>UBB(+1), a mutated form of ubiquitin which is not effectively degraded by the<br>proteasome and is associated with neurogenerative disorders. |
| Cellular Location | Cytoplasm.  |

## Images



Western blot analysis of UCHL3 expression in K562 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.