

GTF2I Antibody

Rabbit mAb Catalog # AP92676

Product Information

Application	WB, IHC, IF, ICC, IHF
Primary Accession	<u>P78347</u>
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	BAP135; BTKAP1; DIWS; Gtf2i; IB291; SPIN; WBS; WBSCR6;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	112416

Additional Information

Dilution Purification Immunogen	WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 Affinity-chromatography A synthesized peptide derived from GTF2I
Description	Interacts with the basal transcription machinery by coordinating the
Storage Condition and Buffer	formation of a multiprotein complex at the C-FOS promoter, and linking specific signal responsive activator complexes. Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	GTF2I
Synonyms	BAP135, WBSCR6
Function	Interacts with the basal transcription machinery by coordinating the formation of a multiprotein complex at the C-FOS promoter, and linking specific signal responsive activator complexes. Promotes the formation of stable high-order complexes of SRF and PHOX1 and interacts cooperatively with PHOX1 to promote serum-inducible transcription of a reporter gene deriven by the C-FOS serum response element (SRE). Acts as a coregulator for USF1 by binding independently two promoter elements, a pyrimidine-rich initiator (Inr) and an upstream E-box. Required for the formation of functional ARID3A DNA- binding complexes and for activation of immunoglobulin heavy-chain transcription upon B-lymphocyte activation.
Cellular Location	Cytoplasm. Nucleus {ECO:0000255 PROSITE-ProRule:PRU00484, ECO:0000269 PubMed:10373551} Note=Colocalizes with BTK in the cytoplasm

Ubiquitous. Isoform 1 is strongly expressed in fetal brain, weakly in adult brain, muscle, and lymphoblasts and is almost undetectable in other adult tissues, while the other isoforms are equally expressed in all adult tissues

Images



Western blot analysis of GTF2I expression in Jurkat cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.