

TDP43 Antibody

Rabbit mAb

Catalog # AP92679

Product Information

Application	WB, IHC, IF, FC, ICC, IHF
Primary Accession	Q13148
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	ALS10; TAR DNA binding protein 43; TARDBP; TDP43;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	44740

Additional Information

Dilution	WB 1:1000~1:5000 IHC 1:50~1:200 ICC/IF 1:50~1:200 FC 1:50
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from TDP43
Description	DNA and RNA-binding protein which regulates transcription and splicing. Involved in the regulation of CFTR splicing. It promotes CFTR exon 9 skipping by binding to the UG repeated motifs in the polymorphic region near the 3'-splice site of this exon. The resulting aberrant splicing is associated with pathological features typical of cystic fibrosis.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	TARDBP {ECO:0000303 PubMed:18396105, ECO:0000312 HGNC:HGNC:11571}
Function	RNA-binding protein that is involved in various steps of RNA biogenesis and processing (PubMed: 23519609). Preferentially binds, via its two RNA recognition motifs RRM1 and RRM2, to GU-repeats on RNA molecules predominantly localized within long introns and in the 3'UTR of mRNAs (PubMed: 23519609 , PubMed: 24240615 , PubMed: 24464995). In turn, regulates the splicing of many non-coding and protein-coding RNAs including proteins involved in neuronal survival, as well as mRNAs that encode proteins relevant for neurodegenerative diseases (PubMed: 21358640 , PubMed: 29438978). Plays a role in maintaining mitochondrial homeostasis by regulating the processing of mitochondrial transcripts (PubMed: 28794432). Also regulates mRNA stability by recruiting CNOT7/CAF1 deadenylase on mRNA 3'UTR leading to poly(A) tail deadenylation and thus shortening (PubMed: 30520513). In response to oxidative insult, associates with stalled

ribosomes localized to stress granules (SGs) and contributes to cell survival (PubMed:[19765185](#), PubMed:[23398327](#)). Also participates in the normal skeletal muscle formation and regeneration, forming cytoplasmic myo-granules and binding mRNAs that encode sarcomeric proteins (PubMed:[30464263](#)). Plays a role in the maintenance of the circadian clock periodicity via stabilization of the CRY1 and CRY2 proteins in a FBXL3-dependent manner (PubMed:[27123980](#)). Negatively regulates the expression of CDK6 (PubMed:[19760257](#)). Regulates the expression of HDAC6, ATG7 and VCP in a PPIA/CYPA-dependent manner (PubMed:[25678563](#)).


Cellular Location

Nucleus. Cytoplasm. Cytoplasm, Stress granule Mitochondrion.
Note=Continuously travels in and out of the nucleus (PubMed:18957508). Localizes to stress granules in response to oxidative stress (PubMed:19765185). A small subset localizes in mitochondria (PubMed:28794432).

Tissue Location

Ubiquitously expressed. In particular, expression is high in pancreas, placenta, lung, genital tract and spleen

Images



Western blot analysis of TDP43 expression in (1) HeLa cell lysate; (2) Mouse brain lysate; (3) Rat brain lysate.

Image not found : 202311/AP92679-IHC.jpg

Immunohistochemical analysis of paraffin-embedded human glioma, using TDP43 Antibody.

Image not found : 202311/AP92679-IF.jpg

Immunofluorescent analysis of Hela cells, using TDP43 Antibody .

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.