

# Erlin-2 Antibody

Rabbit mAb Catalog # AP92700

### **Product Information**

**Application** WB, IHC, IF, ICC, IHF

Primary Accession 094905

Reactivity Human, Mouse Clonality Monoclonal

Other Names ERLIN2; NET32; SPFH2;

IsotypeRabbit IgGHostRabbitCalculated MW37840

#### **Additional Information**

**Dilution** WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200

**Purification** Affinity-chromatography

**Immunogen** A synthesized peptide derived from human Erlin-2

**Description** Component of the ERLIN1/ERLIN2 complex which mediates the endoplasmic

reticulum-associated degradation (ERAD) of inositol 1,4,5-trisphosphate receptors (IP3Rs). Also involved in ITPR1 degradation by the ERAD pathway. Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium

Storage Condition and Buffer

azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term.

Avoid freeze / thaw cycle.

#### **Protein Information**

Name ERLIN2

**Synonyms** C8orf2, SPFH2

**Function** Component of the ERLIN1/ERLIN2 complex which mediates the endoplasmic

reticulum-associated degradation (ERAD) of inositol 1,4,5- trisphosphate receptors (IP3Rs) such as ITPR1 (PubMed:17502376, PubMed:19240031). Promotes sterol-accelerated ERAD of HMGCR probably implicating an AMFR/gp78-containing ubiquitin ligase complex (PubMed:21343306). Involved

in regulation of cellular cholesterol homeostasis by regulation the SREBP signaling pathway. May promote ER retention of the SCAP-SREBF complex

(PubMed:<u>24217618</u>).


**Cellular Location** Endoplasmic reticulum membrane; Single-pass type II membrane protein.

Note=Associated with lipid raft-like domains of the endoplasmic reticulum

membrane

Tissue Location Ubiquitous...

## **Images**



Western blot analysis of Erlin-2 expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.