

Human IgG1 Antibody

Rabbit mAb
Catalog # AP92707

Product Information

Application	WB
Primary Accession	P01857
Reactivity	Human
Clonality	Monoclonal
Other Names	Ig gamma 1 chain C region; IGHG1; Immunoglobulin Gm1;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	43912

Additional Information

Dilution	WB 1:500~1:2000
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human IgG1
Description	There are four IgG subclasses (IgG1, 2, 3 and 4) in humans, named in order of their abundance in serum (IgG1 being the most abundant).
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information


Name	IGHG1 {ECO:0000303 PubMed:11340299, ECO:0000303 Ref.14}
Function	Constant region of immunoglobulin (Ig) heavy chains. Ig are membrane-bound or secreted glycoproteins produced by B lymphocytes. In the recognition phase of humoral immunity, the membrane-bound Ig serve as receptors, which upon binding to a specific antigen trigger the clonal expansion and differentiation of B lymphocytes into Ig-secreting plasma cells. Secreted Ig known as antibodies mediate the effector phase of humoral immunity by blocking the interaction of infectious antigens with cellular receptors (via the antigen-binding region) and eliciting effector mechanisms that lead to pathogen neutralization (via the constant region) (PubMed: 17576170 , PubMed: 20176268 , PubMed: 22158414). The antigen-binding region is formed by the variable domain of one heavy chain paired with the variable domain of its associated light chain. Each Ig molecule has two antigen-binding sites with remarkable affinity for a particular antigen due to V-(D)-J rearrangement, somatic hypermutations and affinity maturation of the variable domains upon antigen exposure (PubMed: 17576170 , PubMed: 20176268 , PubMed: 22158414). The constant region defines the Ig isotype that perform distinct sets of effector functions. B cells diversify and

rearrange their Ig constant regions through class-switch recombination, a process by which the constant region is switched from one Ig isotype to another, namely from IgM and IgD to IgG, IgA and IgE (PubMed:[17576170](#), PubMed:[20176268](#), PubMed:[22158414](#)). The constant region of Ig gamma-1 (IgG1) isotype interacts (via the fragment crystallizable, Fc) with receptors on innate immune cells and the complement system to mediate humoral effector functions, including antibody-dependent cellular cytotoxicity or phagocytosis, complement- dependent cytotoxicity and inflammatory responses.

Cellular Location

[Isoform 1]: Secreted

Images



Western blot analysis of human IgG1 expression in Human tonsil cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.