

# Histone H2B (mono methyl K116) Antibody

Rabbit mAb Catalog # AP92711

### **Product Information**

**Application** WB, IHC, IF, ICC, IHF

Primary Accession Q16778

**Reactivity** Rat, Human, Mouse

**Clonality** Monoclonal **Other Names** Histone H2B;

IsotypeRabbit IgGHostRabbitCalculated MW13920

## **Additional Information**

**Dilution** WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200

**Purification** Affinity-chromatography

ImmunogenA synthesized peptide derived from human Histone H2B (mono methyl K116)DescriptionCore component of nucleosome. Nucleosomes wrap and compact DNA into

chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications

of histones, also called histone code, and nucleosome remodeling.

**Storage Condition and Buffer** Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium

azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term.

Avoid freeze / thaw cycle.

#### **Protein Information**

Name H2BC21 ( <u>HGNC:4760</u>)


**Function** Core component of nucleosome. Nucleosomes wrap and compact DNA into

chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications

of histones, also called histone code, and nucleosome remodeling.

**Cellular Location** Nucleus. Chromosome.

# **Images**



Western blot analysis of Histone H2B (mono methyl K116) expression in Hela cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.