

Fukutin Antibody

Rabbit mAb

Catalog # AP92729

Product Information

Application	WB, IHC
Primary Accession	O75072
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	CMD1X; FCMD; FKTN; Fukutin; LGMD2M; MDDGA4; MDDGB4; MDDGC4;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	53724

Additional Information

Dilution	WB 1:500~1:2000 IHC 1:50~1:200
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Fukutin
Description	May be a glycosyltransferase which participates in glycosylation of alpha-dystroglycan/DAG1. May interact with and reinforce a large complex encompassing the outside and inside of muscle membranes. Could be involved in brain development.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	FKTN (HGNC:3622)
Function	Catalyzes the transfer of a ribitol-phosphate from CDP- ribitol to the distal N-acetylgalactosamine of the phosphorylated O- mannosyl trisaccharide (N-acetylgalactosamine-beta-3-N-acetylglucosamine-beta-4-(phosphate-6-)mannose), a carbohydrate structure present in alpha-dystroglycan (DAG1) (PubMed: 26923585 , PubMed: 27194101 , PubMed: 29477842). This constitutes the first step in the formation of the ribitol 5-phosphate tandem repeat which links the phosphorylated O-mannosyl trisaccharide to the ligand binding moiety composed of repeats of 3-xylosyl-alpha-1,3-glucuronic acid-beta-1 (PubMed: 17034757 , PubMed: 25279699 , PubMed: 26923585 , PubMed: 27194101 , PubMed: 29477842). Required for normal location of POMGNT1 in Golgi membranes, and for normal POMGNT1 activity (PubMed: 17034757). May interact with and reinforce a large complex encompassing the outside and inside of muscle membranes (PubMed: 25279699). Could be involved in brain development (Probable).


Cellular Location

Golgi apparatus membrane; Single-pass type II membrane protein. Cytoplasm {ECO:0000250|UniProtKB:Q8R507}. Nucleus {ECO:0000250|UniProtKB:Q8R507}. Note=In retinal tissue, does not localize with the Golgi apparatus. {ECO:0000250|UniProtKB:Q8R507}

Tissue Location

Expressed in the retina (at protein level) (PubMed:29416295). Widely expressed with highest expression in brain, heart, pancreas and skeletal muscle (PubMed:11115853). Expressed at similar levels in control fetal and adult brain (PubMed:11115853) Expressed in migrating neurons, including Cajal-Retzius cells and adult cortical neurons, as well as hippocampal pyramidal cells and cerebellar Purkinje cells (PubMed:11115853). No expression observed in the glia limitans, the subpial astrocytes (which contribute to basement membrane formation) or other glial cells (PubMed:11115853)

Images



Western blot analysis of Fukutin expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.