

ITPA Antibody

Rabbit mAb

Catalog # AP92771

Product Information

Application	WB, IHC
Primary Accession	Q9BY32
Reactivity	Human
Clonality	Monoclonal
Other Names	Inosine triphosphatase; Itpa; ITPase; My049; NTPase;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	21446


Additional Information

Dilution	WB 1:500~1:2000 IHC 1:50~1:200
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human ITPA
Description	Hydrolyzes ITP and dITP to their respective monophosphate derivatives.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	ITPA {ECO:0000255 HAMAP-Rule:MF_03148}
Synonyms	C20orf37
Function	Pyrophosphatase that hydrolyzes the non-canonical purine nucleotides inosine triphosphate (ITP), deoxyinosine triphosphate (dITP) as well as 2'-deoxy-N-6-hydroxylaminopurine triphosphate (dHAPTP) and xanthosine 5'-triphosphate (XTP) to their respective monophosphate derivatives. The enzyme does not distinguish between the deoxy- and ribose forms. Probably excludes non-canonical purines from RNA and DNA precursor pools, thus preventing their incorporation into RNA and DNA and avoiding chromosomal lesions.
Cellular Location	Cytoplasm {ECO:0000255 HAMAP-Rule:MF_03148, ECO:0000269 PubMed:11278832}
Tissue Location	Ubiquitous. Highly expressed in heart, liver, sex glands, thyroid and adrenal gland

Images



Western blot analysis of ITPA expression in HepG2 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.