10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# ADRA1A Antibody

Rabbit mAb
Catalog # AP92800

## Product Information

| | |
|---|---|
| **Application** | WB, IF, FC, ICC |
| **Primary Accession** | P35348 |
| **Reactivity** | Rat, Human, Mouse |
| **Clonality** | Monoclonal |
| **Other Names** | ADRA1A; Adra1c; ADRA1L1; ALPHA1AAR; |
| **Isotype** | Rabbit IgG |
| **Host** | Rabbit |
| **Calculated MW** | 51487 |

## Additional Information


| | |
|---|---|
| **Dilution** | WB 1:500~1:2000 ICC/IF 1:50~1:200 FC 1:50 |
| **Purification** | Affinity-chromatography |
| **Immunogen** | A synthesized peptide derived from human ADRA1A |
| **Description** | This alpha-adrenergic receptor mediates its action by association with G proteins that activate a phosphatidylinositol-calcium second messenger system. Its effect is mediated by G(q) and G(11) proteins. |
| **Storage Condition and Buffer** | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle. |

## Protein Information

| | |
|---|---|
| **Name** | ADRA1A ( HGNC:277) |
| **Synonyms** | ADRA1C |
| **Function** | Alpha-1 adrenergic receptors are G protein-coupled receptors for catecholamines that signal through the G(q) family of G proteins, including G(q) and G(11). Upon activation, they stimulate the phosphatidylinositol-calcium second messenger pathway, leading to calcium release from intracellular stores and activation of protein kinase C (PubMed:37563160). ADRA1A binds the catecholamine ligands norepinephrine and epinephrine (PubMed:18802028, PubMed:37563160, PubMed:7815325, PubMed:8024574, PubMed:8183249, PubMed:8832064). Can also couple to G(14) protein (By similarity). Nuclear ADRA1A forms heterooligomers with ADRA1B to regulate phenylephrine(PE)-stimulated ERK signaling in cardiac myocytes (PubMed:18802028, PubMed:22120526). At the plasma membrane, ADRA1A interacts with CAVIN4/MURC to regulates ERK activation in cardiomyocytes, contributing to the regulation of cardiac hypertrophy (PubMed:24567387). Additionally, functions as a vasopressor in |

| | |
|---|---|
| | resistance arteries and plays a role in maintaining normal arterial blood pressure (By similarity). |
| **Cellular Location** | Nucleus membrane; Multi-pass membrane protein. Cell membrane; Multi-pass membrane protein. Cytoplasm. Membrane, caveola. Note=Location at the nuclear membrane facilitates heterooligomerization and regulates ERK-mediated signaling in cardiac myocytes (PubMed:18802028, PubMed:22120526) Colocalizes with GNAQ, PLCB1 as well as LAP2 at the nuclear membrane of cardiac myocytes (PubMed:18802028, PubMed:22120526). Colocalizes with CAVIN4 and CAV3 at the plasma membrane and partly within the cytoplasm in cardiomyocytes (PubMed:24567387). |
| **Tissue Location** | Expressed in heart, brain, liver and prostate, but not in kidney, lung, adrenal, aorta and pituitary. Within the prostate, expressed in the apex, base, periurethral and lateral lobe. Isoform 4 is the most abundant isoform expressed in the prostate with high levels also detected in liver and heart. |

## Images




Western blot analysis of ADRA1A expression in HepG2 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.