

p60 katanin Antibody

Rabbit mAb

Catalog # AP92849

Product Information

Application	WB, IF, ICC
Primary Accession	O75449
Reactivity	Human
Clonality	Monoclonal
Other Names	Katanin p60 ATPase containing subunit A1; Katanin p60 subunit A1; KATNA1; p60 katanin;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	55965

Additional Information


Dilution	WB 1:500~1:2000 ICC/IF 1:50~1:200
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human p60 katanin
Description	Severs microtubules in vitro in an ATP-dependent manner. This activity may promote rapid reorganization of cellular microtubule arrays, such as during disassembly of interphase microtubules at the G2-M transition.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	KATNA1 {ECO:0000255 HAMAP-Rule:MF_03023}
Function	Catalytic subunit of a complex which severs microtubules in an ATP-dependent manner. Microtubule severing may promote rapid reorganization of cellular microtubule arrays and the release of microtubules from the centrosome following nucleation. Microtubule release from the mitotic spindle poles may allow depolymerization of the microtubule end proximal to the spindle pole, leading to poleward microtubule flux and poleward motion of chromosome. Microtubule release within the cell body of neurons may be required for their transport into neuronal processes by microtubule-dependent motor proteins. This transport is required for axonal growth.
Cellular Location	Cytoplasm. Midbody. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome {ECO:0000255 HAMAP-Rule:MF_03023} Cytoplasm, cytoskeleton, spindle pole. Cytoplasm, cytoskeleton, spindle. Note=Predominantly cytoplasmic (PubMed:9658175). Localized diffusely in the cytoplasm during

the interphase (PubMed:10751153). During metaphase is localized throughout the cell and more widely dispersed than the microtubules. In anaphase and telophase is localized at the midbody region (PubMed:19261606). Also localized to the interphase centrosome and the mitotic spindle poles (By similarity). Enhanced recruitment to the mitotic spindle poles requires microtubules and interaction with KATNB1 (PubMed:10751153). Localizes within the cytoplasm, partially overlapping with microtubules, in interphase and to the mitotic spindle and spindle poles during mitosis (PubMed:26929214). {ECO:0000255|HAMAP- Rule:MF_03023, ECO:0000269|PubMed:10751153, ECO:0000269|PubMed:19261606, ECO:0000269|PubMed:26929214, ECO:0000269|PubMed:9658175}

Images



Western blot analysis of p60 katanin expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.