

HMG4 Antibody

Rabbit mAb

Catalog # AP92868

Product Information

Application	WB
Primary Accession	O15347
Reactivity	Human, Mouse
Clonality	Monoclonal
Other Names	HMG 4; HMG2A; Hmgb3;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	22980

Additional Information

Dilution	WB 1:500~1:2000
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human HMG4
Description	Binds preferentially single-stranded DNA and unwinds double stranded DNA.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.


Protein Information

Name	HMGB3
Synonyms	HMG2A, HMG4
Function	Multifunctional protein with various roles in different cellular compartments. May act in a redox sensitive manner. Associates with chromatin and binds DNA with a preference for non-canonical DNA structures such as single-stranded DNA. Can bend DNA and enhance DNA flexibility by looping thus providing a mechanism to promote activities on various gene promoters (By similarity). Proposed to be involved in the innate immune response to nucleic acids by acting as a cytoplasmic promiscuous immunogenic DNA/RNA sensor (By similarity). Negatively regulates B-cell and myeloid cell differentiation. In hematopoietic stem cells may regulate the balance between self-renewal and differentiation. Involved in negative regulation of canonical Wnt signaling (By similarity).
Cellular Location	Nucleus {ECO:0000250 UniProtKB:P40618, ECO:0000255 PROSITE-ProRule:PRU00267}. Chromosome Cytoplasm {ECO:0000250 UniProtKB:O54879}

Tissue Location

Expressed predominantly in placenta.

Images



Western blot analysis of HMG4 expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.