

Importin 9 / RANBP9 Antibody

Rabbit mAb

Catalog # AP92894

Product Information

Application	WB, IHC
Primary Accession	Q96P70
Reactivity	Human
Clonality	Monoclonal
Other Names	Imp9; Imp9a; Imp9b; Ipo9; RanBP9;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	115963

Additional Information

Dilution	WB 1:500~1:2000 IHC 1:50~1:200
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Importin 9 / RANBP9
Description	Functions in nuclear protein import as nuclear transport receptor. Serves as receptor for nuclear localization signals (NLS) in cargo substrates. Is thought to mediate docking of the importin/substrate complex to the nuclear pore complex (NPC) through binding to nucleoporin and the complex is subsequently translocated through the pore by an energy requiring, Ran-dependent mechanism.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information


Name	IPO9 {ECO:0000303 PubMed:30855230, ECO:0000312 HGNC:HGNC:19425}
Function	Nuclear transport receptor that mediates nuclear import of proteins, such as histones, proteasome and actin (PubMed: 11823430 , PubMed: 30855230 , PubMed: 34711951). Serves as receptor for nuclear localization signals (NLS) in cargo substrates (PubMed: 11823430). Is thought to mediate docking of the importin/substrate complex to the nuclear pore complex (NPC) through binding to nucleoporin and the complex is subsequently translocated through the pore by an energy requiring, Ran-dependent mechanism (PubMed: 11823430). At the nucleoplasmic side of the NPC, Ran binds to the importin, the importin/substrate complex dissociates and importin is re-exported from the nucleus to the cytoplasm where GTP hydrolysis releases Ran (PubMed: 11823430). The directionality of nuclear import is thought to be conferred by an asymmetric distribution of the GTP- and GDP-bound forms of Ran between the cytoplasm and nucleus (PubMed: 11823430). Mediates the

import of pre-assembled proteasomes into the nucleus; AKIRIN2 acts as a molecular bridge between IPO9 and the proteasome complex (PubMed:[11823430](#), PubMed:[34711951](#)). Mediates the nuclear import of histones H2A, H2B, H4 and H4 (PubMed:[11823430](#), PubMed:[30855230](#)). In addition to nuclear import, also acts as a chaperone for histones by preventing inappropriate non-nucleosomal interactions (PubMed:[30855230](#)). Mediates the nuclear import of actin (By similarity).

Cellular Location

Cytoplasm. Nucleus

Images



Western blot analysis of Importin 9 / RANBP9 expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.