

BCKDK Antibody

Rabbit mAb

Catalog # AP92905

Product Information

Application	WB
Primary Accession	O14874
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	BCKD kinase; BCKDHKIN; Bckdk; BCKDKD; BDK;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	46360

Additional Information

Dilution	WB 1:500~1:2000
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human BCKDK
Description	Catalyzes the phosphorylation and inactivation of the branched-chain alpha-ketoacid dehydrogenase complex, the key regulatory enzyme of the valine, leucine and isoleucine catabolic pathways. Key enzyme that regulate the activity state of the BCKD complex.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	BCKDK {ECO:0000303 PubMed:29779826, ECO:0000312 HGNC:HGNC:16902}
Function	Serine/threonine-protein kinase component of macronutrients metabolism. Forms a functional kinase and phosphatase pair with PPM1K, serving as a metabolic regulatory node that coordinates branched-chain amino acids (BCAAs) with glucose and lipid metabolism via two distinct phosphoprotein targets: mitochondrial BCKDHA subunit of the branched- chain alpha-ketoacid dehydrogenase (BCKDH) complex and cytosolic ACLY, a lipogenic enzyme of Krebs cycle (PubMed: 24449431 , PubMed: 29779826 , PubMed: 37558654). Phosphorylates and inactivates mitochondrial BCKDH complex a multisubunit complex consisting of three multimeric components each involved in different steps of BCAA catabolism: E1 composed of BCKDHA and BCKDHB, E2 core composed of DBT monomers, and E3 composed of DLD monomers. Associates with the E2 component of BCKDH complex and phosphorylates BCKDHA on Ser-337, leading to conformational changes that interrupt substrate channeling between E1 and E2 and inactivates the BCKDH complex

(PubMed:[29779826](#), PubMed:[37558654](#)). Phosphorylates ACLY on Ser-455 in response to changes in cellular carbohydrate abundance such as occurs during fasting to feeding metabolic transition. Refeeding stimulates MLXIPL/ChREBP transcription factor, leading to increased BCKDK to PPM1K expression ratio, phosphorylation and activation of ACLY that ultimately results in the generation of malonyl-CoA and oxaloacetate immediate substrates of de novo lipogenesis and gluconeogenesis, respectively (PubMed:[29779826](#)). Recognizes phosphosites having SxxE/D canonical motif (PubMed:[29779826](#)).


Cellular Location

Mitochondrion matrix {ECO:0000250 | UniProtKB:Q00972, ECO:0000305 | PubMed:24449431} Note=Detected in the cytosolic compartment of liver cells {ECO:0000250 | UniProtKB:Q00972}

Tissue Location

Ubiquitous.

Images



Western blot analysis of BCKDK expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.