

# NUP50 Antibody

Rabbit mAb

Catalog # AP92924

## Product Information

---

<b>Application</b>	WB, IHC, IF, ICC, IHF
<b>Primary Accession</b>	<a href="#">Q9UKX7</a>
<b>Reactivity</b>	Rat, Human, Mouse
<b>Clonality</b>	Monoclonal
<b>Other Names</b>	NPAP60; NPAP60L; Nup50;
<b>Isotype</b>	Rabbit IgG
<b>Host</b>	Rabbit
<b>Calculated MW</b>	50144

## Additional Information

---

<b>Dilution</b>	WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200
<b>Purification</b>	Affinity-chromatography
<b>Immunogen</b>	A synthesized peptide derived from human NUP50
<b>Description</b>	Component of the nuclear pore complex that has a direct role in nuclear protein import. Actively displaces NLSs from importin-alpha, and facilitates disassembly of the importin-alpha:beta-cargo complex and importin recycling.
<b>Storage Condition and Buffer</b>	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

## Protein Information

---

<b>Name</b>	NUP50
<b>Synonyms</b>	NPAP60L
<b>Function</b>	Component of the nuclear pore complex that has a direct role in nuclear protein import (PubMed: <a href="#">20016008</a> ). Actively displaces NLSs from importin-alpha, and facilitates disassembly of the importin- alpha:beta-cargo complex and importin recycling (PubMed: <a href="#">20016008</a> ). Interacts with regulatory proteins of cell cycle progression including CDKN1B (By similarity). This interaction is required for correct intracellular transport and degradation of CDKN1B (By similarity).
<b>Cellular Location</b>	Nucleus, nuclear pore complex. Nucleus membrane {ECO:0000250 UniProtKB:O08587}; Peripheral membrane protein {ECO:0000250 UniProtKB:O08587}; Nucleoplasmic side {ECO:0000250 UniProtKB:O08587}. Note=Localizes to the nucleoplasmic fibrils of the nuclear pore complex (By similarity). Dissociates from the NPC


structure early during prophase of mitosis (PubMed:12802065) Associates with the newly formed nuclear membrane during telophase (PubMed:12802065). In the testis, the localization changes during germ cell differentiation from the nuclear surface in spermatocytes to the whole nucleus (interior) in spermatids and back to the nuclear surface in spermatozoa (By similarity). {ECO:0000250|UniProtKB:O08587, ECO:0000269|PubMed:12802065}

#### Tissue Location

Ubiquitous. Highest levels in testis, peripheral blood leukocytes and fetal liver

## Images

---



Western blot analysis of NUP50 expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.