

GNE Antibody

Rabbit mAb

Catalog # AP93000

Product Information

Application	WB, IF, ICC
Primary Accession	Q9Y223
Reactivity	Human
Clonality	Monoclonal
Other Names	DMRV; GNE; IBM2; ManAc kinase; Uae1;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	79275

Additional Information

Dilution	WB 1:500~1:2000 ICC/IF 1:50~1:200
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human GNE
Description	Regulates and initiates biosynthesis of N-acetylneuraminic acid (NeuAc), a precursor of sialic acids. Plays an essential role in early development (By similarity). Required for normal sialylation in hematopoietic cells. Sialylation is implicated in cell adhesion, signal transduction, tumorigenicity and metastatic behavior of malignant cells.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	GNE (HGNC:23657)
Function	Bifunctional enzyme that possesses both UDP-N- acetylglucosamine 2-epimerase and N-acetylmannosamine kinase activities, and serves as the initiator of the biosynthetic pathway leading to the production of N-acetylneuraminic acid (NeuAc), a critical precursor in the synthesis of sialic acids. By catalyzing this pivotal and rate-limiting step in sialic acid biosynthesis, this enzyme assumes a pivotal role in governing the regulation of cell surface sialylation, playing a role in embryonic angiogenesis (PubMed: 10334995 , PubMed: 11326336 , PubMed: 14707127 , PubMed: 16503651 , PubMed: 2808337 , PubMed: 38237079). Sialic acids represent a category of negatively charged sugars that reside on the surface of cells as terminal components of glycoconjugates and mediate important functions in various cellular processes, including cell adhesion, signal transduction, and cellular recognition (PubMed: 10334995 , PubMed: 14707127).


Cellular Location

Cytoplasm, cytosol {ECO:0000250|UniProtKB:O35826}

Tissue Location

Highest expression in liver and placenta. Also found in heart, brain, lung, kidney, skeletal muscle and pancreas Isoform 1 is expressed in heart, brain, kidney, liver, placenta, lung, spleen, pancreas, skeletal muscle and colon. Isoform 2 is expressed mainly in placenta, but also in brain, kidney, liver, lung, pancreas and colon. Isoform 3 is expressed at low level in kidney, liver, placenta and colon.

Images



Western blot analysis of GNE expression in K562 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.