10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Phospho-PBK/TOPK (Thr9) Antibody

Rabbit mAb
Catalog # AP93037

Product Information

Application	WB, IHC
Primary Accession	Q96KB5
Reactivity	Human
Clonality	Monoclonal
Other Names	CT84; MAPKK like protein kinase; Nori3; PBK; PDZ binding kinas; Serine/threonine protein kinase; SPK; TOPK;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	36085


Additional Information

Dilution	WB 1:500~1:2000 IHC 1:50~1:200
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Phospho-PBK/TOPK (Thr9)
Description	Phosphorylates MAP kinase p38. Seems to be active only in mitosis. May also play a role in the activation of lymphoid cells. When phosphorylated, forms a complex with TP53, leading to TP53 destabilization and attenuation of G2/M checkpoint during doxorubicin-induced DNA damage.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	PBK
Synonyms	TOPK
Function	Phosphorylates MAP kinase p38. Seems to be active only in mitosis. May also play a role in the activation of lymphoid cells. When phosphorylated, forms a complex with TP53, leading to TP53 destabilization and attenuation of G2/M checkpoint during doxorubicin- induced DNA damage.
Tissue Location	Expressed in the testis and placenta. In the testis, restrictedly expressed in outer cell layer of seminiferous tubules.

Images



Western blot analysis of Phospho-PBK/TOPK (Thr9) expression in (1) HeLa cell lysate; (2) HeLa cell treated with Nocodazole lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.