

# Phospho-PBK/TOPK (Thr9) Antibody

Rabbit mAb

Catalog # AP93037

## Product Information

---

<b>Application</b>	WB, IHC
<b>Primary Accession</b>	<a href="#">Q96KB5</a>
<b>Reactivity</b>	Human
<b>Clonality</b>	Monoclonal
<b>Other Names</b>	CT84; MAPKK like protein kinase; Nori3; PBK; PDZ binding kinase; Serine/threonine protein kinase; SPK; TOPK;
<b>Isotype</b>	Rabbit IgG
<b>Host</b>	Rabbit
<b>Calculated MW</b>	36085

## Additional Information

---

<b>Dilution</b>	WB 1:500~1:2000 IHC 1:50~1:200
<b>Purification</b>	Affinity-chromatography
<b>Immunogen</b>	A synthesized peptide derived from human Phospho-PBK/TOPK (Thr9)
<b>Description</b>	Phosphorylates MAP kinase p38. Seems to be active only in mitosis. May also play a role in the activation of lymphoid cells. When phosphorylated, forms a complex with TP53, leading to TP53 destabilization and attenuation of G2/M checkpoint during doxorubicin-induced DNA damage.
<b>Storage Condition and Buffer</b>	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.


## Protein Information

---

<b>Name</b>	PBK
<b>Synonyms</b>	TOPK
<b>Function</b>	Phosphorylates MAP kinase p38. Seems to be active only in mitosis. May also play a role in the activation of lymphoid cells. When phosphorylated, forms a complex with TP53, leading to TP53 destabilization and attenuation of G2/M checkpoint during doxorubicin-induced DNA damage.
<b>Tissue Location</b>	Expressed in the testis and placenta. In the testis, restrictedly expressed in outer cell layer of seminiferous tubules.

## Images

---



Western blot analysis of Phospho-PBK/TOPK (Thr9) expression in (1) HeLa cell lysate; (2) HeLa cell treated with Nocodazole lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.