

RNF7 Antibody

Rabbit mAb

Catalog # AP93095

Product Information

Application	WB, IHC, IF, ICC, IHF
Primary Accession	Q9UBF6
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	CKBBP1; Rbx2; RNF7; ROC2; SAG;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	12683

Additional Information

Dilution	WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human RNF7
Description	Probable component of the SCF (SKP1-CUL1-F-box protein) E3 ubiquitin ligase complex which mediates the ubiquitination and subsequent proteasomal degradation of target proteins involved in cell cycle progression, signal transduction and transcription.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	RNF7 (HGNC:10070)
Function	Catalytic component of multiple cullin-5-RING E3 ubiquitin- protein ligase complexes (ECS complexes), which mediate the ubiquitination and subsequent proteasomal degradation of target proteins (PubMed: 21980433 , PubMed: 33268465 , PubMed: 38418882 , PubMed: 38574733 , PubMed: 35512830 , PubMed: 40440427). It is thereby involved in various biological processes, such as cell cycle progression, signal transduction and transcription (PubMed: 21980433 , PubMed: 33268465 , PubMed: 38418882 , PubMed: 38574733). The functional specificity of the E3 ubiquitin-protein ligase ECS complexes depend on the variable SOCS box-containing substrate recognition component (PubMed: 21980433 , PubMed: 33268465). Within ECS complexes, RNF7/RBX2 recruits the E2 ubiquitination enzyme to the complex via its RING-type and brings it into close proximity to the substrate (PubMed: 34518685). Catalytic subunit of various SOCS-containing ECS complexes, such as the ECS(SOCS7) complex, that regulate reelin signaling by mediating ubiquitination and degradation of DAB1 (By similarity). The

ECS(SOCS2) complex mediates the ubiquitination and subsequent proteasomal degradation of phosphorylated EPOR and GHR (PubMed:[21980433](#), PubMed:[25505247](#)). Promotes ubiquitination and degradation of NF1, thereby regulating Ras protein signal transduction (By similarity). As part of the ECS(ASB9) complex, catalyzes ubiquitination and degradation of CKB (PubMed:[33268465](#)). The ECS(SPSB3) complex catalyzes ubiquitination of nuclear CGAS (PubMed:[3841882](#)). As part of the ECS(RAB40C) complex, mediates ANKRD28 ubiquitination and degradation, thereby inhibiting protein phosphatase 6 (PP6) complex activity and focal adhesion assembly during cell migration (PubMed:[35512830](#)). The ECS(ASB7) complex acts a negative regulator of H3K9me3 histone mark by mediating ubiquitination and degradation of SUV39H1 (PubMed:[40440427](#)). As part of some ECS complex, catalyzes 'Lys-11'-linked ubiquitination and degradation of BTRC (PubMed:[27910872](#)). ECS complexes and ARIH2 collaborate in tandem to mediate ubiquitination of target proteins; ARIH2 mediating addition of the first ubiquitin on CRLs targets (PubMed:[34518685](#), PubMed:[3841882](#)). Specifically catalyzes the neddylation of CUL5 via its interaction with UBE2F (PubMed:[19250909](#)). Does not catalyze neddylation of other cullins (CUL1, CUL2, CUL3, CUL4A or CUL4B) (PubMed:[19250909](#)). May play a role in protecting cells from apoptosis induced by redox agents (PubMed:[10082581](#)).


Cellular Location

Cytoplasm. Nucleus

Tissue Location

Expressed in heart, liver, skeletal muscle and pancreas. At very low levels expressed in brain, placenta and lung

Images



Western blot analysis of RNF7 expression in HepG2 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.