

BIN1 Antibody

Rabbit mAb

Catalog # AP93097

Product Information

Application	WB, IHC, IF, FC, ICC, IP, IHF
Primary Accession	O00499
Reactivity	Human
Clonality	Monoclonal
Other Names	AMPH2; Amphiphysin 2; Amphiphysin II; AMPHL; Bin1; Bridging integrator 1; SH3P9;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	64699

Additional Information

Dilution	WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 IP 1:50 FC 1:50
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human BIN1
Description	May be involved in regulation of synaptic vesicle endocytosis. May act as a tumor suppressor and inhibits malignant cell transformation.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information


Name	BIN1
Synonyms	AMPHL
Function	Is a key player in the control of plasma membrane curvature, membrane shaping and membrane remodeling. Required in muscle cells for the formation of T-tubules, tubular invaginations of the plasma membrane that function in depolarization-contraction coupling (PubMed: 24755653). Is a negative regulator of endocytosis (By similarity). Is also involved in the regulation of intracellular vesicles sorting, modulation of BACE1 trafficking and the control of amyloid-beta production (PubMed: 27179792). In neuronal circuits, endocytosis regulation may influence the internalization of PHF-tau aggregates (By similarity). May be involved in the regulation of MYC activity and the control cell proliferation (PubMed: 8782822). Has actin bundling activity and stabilizes actin filaments against depolymerization in vitro (PubMed: 28893863).
Cellular Location	[Isoform BIN1]: Nucleus. Cytoplasm Endosome

{ECO:0000250|UniProtKB:O08539}. Cell membrane, sarcolemma, T- tubule
{ECO:0000250|UniProtKB:O08839}

Tissue Location

Ubiquitous. Highest expression in the brain and muscle (PubMed:9182667). Expressed in oligodendrocytes (PubMed:27488240). Isoform IIA is expressed only in the brain, where it is detected in the gray matter, but not in the white matter (PubMed:27488240). Isoform BIN1 is widely expressed with highest expression in skeletal muscle.

Images



Western blot analysis of BIN1 expression in A431 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.