

Caspase 3 Antibody

Rabbit mAb Catalog # AP93100

Product Information

Application Primary Accession Reactivity Clonality Other Names	WB, IHC, IP <u>P42574</u> Rat, Human, Mouse Monoclonal Apopain precursor; CASP-3; CPP-32; CPP32; Caspase-3; Cysteine protease CPP32; ICE3; SCA-1; SREBP cleavage activity 1;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	31608

Additional Information

Dilution Purification Immunogen Description	WB 1:500~1:2000 IHC 1:50~1:200 IP 1:50 Affinity-chromatography A synthesized peptide derived from human caspase 3 This gene encodes a protein which is a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce 2 subunits, large and small, that dimerize to form the active enzyme
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	CASP3
Synonyms	CPP32 {ECO:0000303 PubMed:7983002}
Function	Thiol protease that acts as a major effector caspase involved in the execution phase of apoptosis (PubMed: <u>18723680</u> , PubMed: <u>20566630</u> , PubMed: <u>23650375</u> , PubMed: <u>35338844</u> , PubMed: <u>35446120</u> , PubMed: <u>7596430</u>). Following cleavage and activation by initiator caspases (CASP8, CASP9 and/or CASP10), mediates execution of apoptosis by catalyzing cleavage of many proteins (PubMed: <u>18723680</u> , PubMed: <u>20566630</u> , PubMed: <u>23650375</u> , PubMed: <u>7596430</u>). At the onset of apoptosis, it proteolytically cleaves poly(ADP-ribose) polymerase PARP1 at a '216-Asp- -Gly-217' bond (PubMed: <u>10497198</u> , PubMed: <u>16374543</u> , PubMed: <u>7596430</u> , PubMed: <u>7596430</u> , PubMed: <u>7596430</u> , PubMed: <u>16374543</u> ,

	element binding proteins (SREBPs) between the basic helix-loop-helix leucine zipper domain and the membrane attachment domain (By similarity). Cleaves and activates caspase-6, -7 and -9 (CASP6, CASP7 and CASP9, respectively) (PubMed: <u>7596430</u>). Cleaves and inactivates interleukin-18 (IL18) (PubMed: <u>37993714</u> , PubMed: <u>9334240</u>). Involved in the cleavage of huntingtin (PubMed: <u>8696339</u>). Triggers cell adhesion in sympathetic neurons through RET cleavage (PubMed: <u>21357690</u>). Cleaves and inhibits serine/threonine-protein kinase AKT1 in response to oxidative stress (PubMed: <u>23152800</u>). Acts as an inhibitor of type I interferon production during virus-induced apoptosis by mediating cleavage of antiviral proteins CGAS, IRF3 and MAVS, thereby preventing cytokine overproduction (PubMed: <u>30878284</u>). Also involved in pyroptosis by mediating cleavage and activation of gasdermin-E (GSDME) (PubMed: <u>35338844</u> , PubMed: <u>35446120</u>). Cleaves XRCC4 and phospholipid scramblase proteins XKR4, XKR8 and XKR9, leading to promote phosphatidylserine exposure on apoptotic cell surface (PubMed: <u>23845944</u> , PubMed: <u>33725486</u>). Cleaves BIRC6 following inhibition of BIRC6-caspase binding by DIABLO/SMAC (PubMed: <u>36758104</u> , PubMed: <u>36758106</u>).
Cellular Location	Cytoplasm.
Tissue Location	Highly expressed in lung, spleen, heart, liver and kidney. Moderate levels in brain and skeletal muscle, and low in testis. Also found in many cell lines, highest expression in cells of the immune system.

Images



Western blot analysis of Calreticulin expression in (1) Jurkat cell lysate; (2) NIH/3T3 cell lysate; (3) Rat brain lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.