

IL22 Antibody

Rabbit mAb Catalog # AP93113

Product Information

Application WB
Primary Accession Q9GZX6
Reactivity Human
Clonality Monoclonal

Other Names IL21; Il22; ILD110; ILTIF; interleukin 21; Interleukin 22; TIFa; TIFIL23; zcyto18;

IsotypeRabbit IgGHostRabbitCalculated MW20011

Additional Information

Dilution WB 1:500~1:2000 **Purification** Affinity-chromatography

Immunogen A synthesized peptide derived from human IL22

Description Cytokine that contributes to the inflammatory response in vivo.

Storage Condition and Buffer Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium

azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term.

Avoid freeze / thaw cycle.

Protein Information

Name IL22

Synonyms ILTIF, ZCYTO18

Function Cytokine that plays a critical role in modulating tissue responses during

inflammation (PubMed:<u>17204547</u>). Plays an essential role in the regeneration of epithelial cells to maintain barrier function after injury and for the prevention of further tissue damage (PubMed:<u>17204547</u>). Unlike most of the cytokines, has no effect on immune cells. Signals through a heterodimeric

receptor composed of two subunits, the specific receptor IL22RA1 which is present on non-immune cells in many organs and the shared subunit IL10RB (PubMed:10875937, PubMed:18599299). Ligation of IL22RA1 with IL22 induces activation of the tyrosine kinases JAK1 and TYK2, which in turn activates STAT3. In turn, promotes cell survival and proliferation through


STAT3, ERK1/2 and PI3K/AKT pathways (PubMed: 25793261,

PubMed:31311100). Promotes phosphorylation of GSK3B at 'Ser-9' and CTTN

(By similarity). Promotes epithelial cell spreading (By similarity).

Cellular Location Secreted.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.