10320 Camino Santa Fe, Suite G San Diego, CA 92121 Tel: 858.875.1900 Fax: 858.875.1999



# **IL9 Antibody**

Rabbit mAb Catalog # AP93114

### **Product Information**

Application WB
Primary Accession P15248
Reactivity Human
Clonality Monoclonal

Other Names HP40; Il9; Interleukin 9 MCGF; P40; TCGF 3;

IsotypeRabbit IgGHostRabbitCalculated MW15909

## **Additional Information**

**Dilution** WB 1:500~1:2000 **Purification** Affinity-chromatography

**Immunogen** A synthesized peptide derived from human IL9

**Description**Supports IL-2 independent and IL-4 independent growth of helper T-cells. **Storage Condition and Buffer**Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium

azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term.

Avoid freeze / thaw cycle.

#### **Protein Information**

Name IL9

**Function** Multifunctional cytokine secreted mainly by T-helper 2 lymphocytes and also

mast cells or NKT cells that plays important roles in the immune response

against parasites (PubMed: 29742432). Affects intestinal epithelial


permeability and adaptive immunity (PubMed: <u>29742432</u>). In addition, induces the differentiation of specific T-cell subsets such as IL-17 producing helper T-cells (TH17) and also proliferation and differentiation of mast cells. Mechanistically, exerts its biological effects through a receptor composed of

IL9R subunit and a signal transducing subunit IL2RG. Receptor stimulation results in the rapid activation of JAK1 and JAK3 kinase activities leading to STAT1, STAT3 and STAT5-mediated transcriptional programs. Induction of differentiation genes seems to be mediated by STAT1 alone, while protection

of cells from apoptosis depends on STAT3 and STAT5.

**Cellular Location** Secreted.

# **Images**



Western blot analysis of IL9 in Recombinant Human IL9 protein cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.