

CD125 Antibody

Rabbit mAb Catalog # AP93118

Product Information

Application WB
Primary Accession Q01344
Reactivity Human
Clonality Monoclonal

Other Names CD125; CDw125; HSIL5R3; IL5R; IL5RA;

IsotypeRabbit IgGHostRabbitCalculated MW47685

Additional Information

Dilution WB 1:500~1:2000 **Purification** Affinity-chromatography

Immunogen A synthesized peptide derived from human CD125

Description This is the receptor for interleukin-5. The alpha chain binds to IL5.

Storage Condition and Buffer Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium

azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term.

Avoid freeze / thaw cycle.

Protein Information

Name IL5RA

Synonyms IL5R

Function Cell surface receptor that plays an important role in the survival,

differentiation, and chemotaxis of eosinophils (PubMed:<u>9378992</u>). Acts by forming a heterodimeric receptor with CSF2RB subunit and subsequently binding to interleukin-5 (PubMed:<u>1495999</u>, PubMed:<u>22528658</u>). In unstimulated conditions, interacts constitutively with JAK2. Heterodimeric receptor activation leads to JAK2 stimulation and subsequent activation of the


Teceptor activation reads to joine still addition and subsequent

JAK-STAT pathway (PubMed: 9516124).

Cellular Location Membrane; Single-pass type I membrane protein.

Tissue Location Expressed on eosinophils and basophils.

Images



Western blot analysis of CD125 in Human serum cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.