10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Phospho-ErbB2(Y1221 + Y1222) Antibody

Rabbit mAb
Catalog # AP93147

Product Information

Application	WB
Primary Accession	P04626
Reactivity	Human
Clonality	Monoclonal
Other Names	CD340; CerbB2; Erb b2 receptor tyrosine kinase 2; ERBB2; HER2; Herstatin; Human epidermal growth factor receptor 2; MLN19; NEU; NGL; Proto-oncogene Neu; Receptor tyrosine-protein kinase erbB-2; Tyrosine kinase type cell surface receptor HER2; V erb b2 avian erythroblastic leukemia viral oncogene homolog 2; V erb b2 avian erythroblastic leukemia viral oncoprotein 2;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	137910

Additional Information


Dilution	WB 1:500~1:2000
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Phospho-ErbB2(Y1221 + Y1222)
Description	Protein tyrosine kinase that is part of several cell surface receptor complexes, but that apparently needs a coreceptor for ligand binding. Essential component of a neuregulin-receptor complex, although neuregulins do not interact with it alone. GP30 is a potential ligand for this receptor.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	ERBB2
Synonyms	HER2, MLN19, NEU, NGL
Function	Protein tyrosine kinase that is part of several cell surface receptor complexes, but that apparently needs a coreceptor for ligand binding. Essential component of a neuregulin-receptor complex, although neuregulins do not interact with it alone. GP30 is a potential ligand for this receptor. Regulates outgrowth and stabilization of peripheral microtubules (MTs). Upon ERBB2 activation, the MEMO1-RHOA-DIAPH1 signaling pathway elicits the phosphorylation and thus the inhibition of GSK3B at cell membrane. This prevents the phosphorylation of APC and CLASP2, allowing its association

	with the cell membrane. In turn, membrane-bound APC allows the localization of MACF1 to the cell membrane, which is required for microtubule capture and stabilization.
Cellular Location	Cell membrane; Single-pass type I membrane protein. Cell projection, ruffle membrane; Single-pass type I membrane protein. Note=Internalized from the cell membrane in response to EGF stimulation. [Isoform 2]: Cytoplasm. Nucleus.
Tissue Location	Expressed in a variety of tumor tissues including primary breast tumors and tumors from small bowel, esophagus, kidney and mouth.

Images



Western blot analysis of Phospho-ErbB2(Y1221 + Y1222) expression in SKBR3 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.