

# Phospho-EGFR (Y869) Antibody

Rabbit mAb Catalog # AP93149

### **Product Information**

Application WB, IF, ICC
Primary Accession P00533
Reactivity Human
Clonality Monoclonal

Other Names EGFR; ERBB; ERBB1; Errp; HER1; mENA; NISBD2; PIG61; SA7; wa2; Wa5;

IsotypeRabbit IgGHostRabbitCalculated MW134277

## **Additional Information**

**Dilution** WB 1:500~1:2000 ICC/IF 1:50~1:200

**Purification** Affinity-chromatography

Immunogen A synthesized peptide derived from human EGFR Full-length sequence 1210aa

around the phosphorylation site of Tyrosine 869

**Description** Receptor tyrosine kinase binding ligands of the EGF family and activating

several signaling cascades to convert extracellular cues into appropriate

cellular responses.

Storage Condition and Buffer Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium

azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term.

Avoid freeze / thaw cycle.

#### **Protein Information**

Name EGFR ( HGNC:3236)

**Synonyms** ERBB, ERBB1, HER1

**Function** Receptor tyrosine kinase binding ligands of the EGF family and activating

several signaling cascades to convert extracellular cues into appropriate cellular responses (PubMed:10805725, PubMed:27153536, PubMed:2790960, PubMed:35538033). Known ligands include EGF, TGFA/TGF- alpha, AREG, epigen/EPGN, BTC/betacellulin, epiregulin/EREG and HBEGF/heparin-binding

EGF (PubMed: 12297049, PubMed: 15611079, PubMed: 17909029,

PubMed: 20837704, PubMed: 27153536, PubMed: 2790960, PubMed: 7679104, PubMed: 8144591, PubMed: 9419975). Ligand binding triggers receptor homoand/or heterodimerization and autophosphorylation on key cytoplasmic residues. The phosphorylated receptor recruits adapter proteins like GRB2 which in turn activates complex downstream signaling cascades. Activates at least 4 major downstream signaling cascades including the RAS-RAF-MEK-ERK, PI3 kinase-AKT, PLCgamma-PKC and STATs modules (PubMed: 27153536). May

also activate the NF-kappa-B signaling cascade (PubMed:11116146). Also directly phosphorylates other proteins like RGS16, activating its GTPase activity and probably coupling the EGF receptor signaling to the G protein-coupled receptor signaling (PubMed:11602604). Also phosphorylates MUC1 and increases its interaction with SRC and CTNNB1/beta-catenin (PubMed:11483589). Positively regulates cell migration via interaction with CCDC88A/GIV which retains EGFR at the cell membrane following ligand stimulation, promoting EGFR signaling which triggers cell migration (PubMed:20462955). Plays a role in enhancing learning and memory performance (By similarity). Plays a role in mammalian pain signaling (long-lasting hypersensitivity) (By similarity).


#### **Cellular Location**

Cell membrane; Single-pass type I membrane protein. Endoplasmic reticulum membrane; Single-pass type I membrane protein Golgi apparatus membrane; Single-pass type I membrane protein. Nucleus membrane; Single-pass type I membrane protein. Endosome. Endosome membrane. Nucleus. Note=In response to EGF, translocated from the cell membrane to the nucleus via Golgi and ER (PubMed:17909029, PubMed:20674546). Endocytosed upon activation by ligand (PubMed:17182860, PubMed:17909029, PubMed:27153536, PubMed:2790960). Colocalized with GPER1 in the nucleus of estrogen agonist-induced cancer-associated fibroblasts (CAF) (PubMed:20551055)

#### **Tissue Location**

Ubiquitously expressed. Isoform 2 is also expressed in ovarian cancers.

## **Images**



Western blot analysis of Phospho-EGFR (Y845) expression in A431 treated with EGF cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.