

Histone H2B (acetyl K16) Antibody

Rabbit mAb Catalog # AP93200

Product Information

Application	WB, IHC, IF, ICC, IP, IHF
Primary Accession	<u>Q16778</u>
Reactivity	Human, Mouse
Clonality	Monoclonal
Other Names	Histone H2B type 1-H; Histone H2B.j; H2B/j; HIST1H2BH; H2BFJ;
lsotype	Rabbit IgG
Host	Rabbit
Calculated MW	13920


Additional Information

Dilution Purification Immunogen	WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 Affinity-chromatography A synthesized peptide derived from human Histone H2B (acetyl K16)
Description	Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	H2BC21 (<u>HGNC:4760</u>)
Function	Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.
Cellular Location	Nucleus. Chromosome.
Images	

Western blot analysis of Histone H2B (acetyl K16) expression in HeLa treated with Trichostatin A cell lysate.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.