

SDHC Antibody

Rabbit mAb Catalog # AP93236

Product Information

Application WB, IHC Primary Accession Q99643

Reactivity Rat, Human, Mouse

Clonality Monoclonal

Other Names CYBL; PGL3; QPs1; SDH3; sdhC;

IsotypeRabbit IgGHostRabbitCalculated MW18610

Additional Information

Dilution WB 1:500~1:2000 IHC 1:50~1:200

Purification Affinity-chromatography

Immunogen A synthesized peptide derived from human SDHC

DescriptionMembrane-anchoring subunit of succinate dehydrogenase (SDH) that is involved in complex II of the mitochondrial electron transport chain and is

involved in complex II of the mitochondrial electron transport chain and is responsible for transferring electrons from succinate to ubiquinone

(coenzyme Q).

Storage Condition and Buffer Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium

azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term.

Avoid freeze / thaw cycle.

Protein Information

Name SDHC

Synonyms CYB560, SDH3

Function Membrane-anchoring subunit of succinate dehydrogenase (SDH) that is

involved in complex II of the mitochondrial electron transport chain and is responsible for transferring electrons from succinate to ubiquinone (coenzyme Q) (PubMed:9533030). SDH also oxidizes malate to the

non-canonical enol form of oxaloacetate, enol-oxaloacetate (By similarity). Enol-oxaloacetate, which is a potent inhibitor of the succinate dehydrogenase

activity, is further isomerized into keto- oxaloacetate (By similarity).

Cellular Location Mitochondrion inner membrane; Multi-pass membrane protein

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.