

RPLP2 Antibody (N-Term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP9327a

Product Information

Application	WB, E
Primary Accession	P05387
Other Accession	P19943 , P99027 , P42899 , NP_000995
Reactivity	Human, Rat, Mouse
Predicted	Bovine, Mouse, Rabbit
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB40737
Calculated MW	11665
Antigen Region	13-41

Additional Information

Gene ID	6181
Other Names	60S acidic ribosomal protein P2, Renal carcinoma antigen NY-REN-44, RPLP2, D11S2243E, RPP2
Target/Specificity	This RPLP2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 13-41 amino acids from the N-terminal region of human RPLP2.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	RPLP2 Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	RPLP2
Synonyms	D11S2243E, RPP2

Function

Plays an important role in the elongation step of protein synthesis.


Background

RPLP2 consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This protein encodes a ribosomal phosphoprotein that is a component of the 60S subunit. The protein, which is a functional equivalent of the E. coli L7/L12 ribosomal protein, belongs to the L12P family of ribosomal proteins. It plays an important role in the elongation step of protein synthesis. Unlike most ribosomal proteins, which are basic, the encoded protein is acidic. Its C-terminal end is nearly identical to the C-terminal ends of the ribosomal phosphoproteins P0 and P1. The P2 protein can interact with P0 and P1 to form a pentameric complex consisting of P1 and P2 dimers, and a P0 monomer. The protein is located in the cytoplasm. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome.

References

Martinez-Azorin,F. FEBS Lett. 582 (20), 3029-3032 (2008) Martinez-Azorin,F. Biochem. J. 413 (3), 527-534 (2008) Sugiyama,N. Mol. Cell Proteomics 6 (6), 1103-1109 (2007)

Images



RPLP2 Antibody (N-Term) (Cat. #AP9327a) western blot analysis in K562 cell line lysates (35ug/lane). This demonstrates the RPLP2 antibody detected the RPLP2 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.