10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-FABP6 Antibody

Catalog # AP93283

Product Information

Application	WB, IF/IC
Primary Accession	P51161
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	14371

Additional Information


Gene ID	2172
Other Names	ILBP; ILLBP; Gastrotropin; GT; Fatty acid-binding protein 6; Ileal lipid-binding protein; ILBP; Intestinal 15 kDa protein; I-15P; Intestinal bile acid-binding protein; I-BABP
Target/Specificity	Recombinant fusion protein of human FABP6
Dilution	WB~~WB (1/500 - 1/1000) IF/IC~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	FABP6
Synonyms	ILBP, ILLBP
Function	Binds to bile acids and is involved in enterohepatic bile acid metabolism. Required for efficient apical to basolateral transport of conjugated bile acids in ileal enterocytes (By similarity). In vitro binds to bile acids in the order: deoxycholic acid > cholic acid > chenodeoxycholic acid and respective BA conjugation modifies affinities in the order taurine-conjugated > glycine-conjugated > unconjugated bile acids. Stimulates gastric acid and pepsinogen secretion (By similarity).
Cellular Location	[Isoform 1]: Cytoplasm {ECO:0000250|UniProtKB:P80020}. Membrane; Peripheral membrane protein {ECO:0000250|UniProtKB:P50119}; Cytoplasmic side {ECO:0000250|UniProtKB:P50119}
Tissue Location	Isoform 1 is expressed in the jejunum, ileum, cecum and ascending colon

intestine. Isoform 2 is xpressed in the gallbladder, duodenum, jejunum, ileum, cecum, ascending, transverse and descending colon, sigmoid colon and rectum. Isoform 2 is expressed in colorectal adenocarcinomas and their adjacent normal mucosa (at protein level).

Images



Western blot analysis of FABP6 expression in MCF7 (A), A375 (B) whole cell lysates. (Predicted band size: 14 kD; Observed band size: 18 kD)



Immunofluorescent analysis of FABP6 staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.