

DLL3 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP9328B

Product Information

Application	IHC-P-Leica, IF, WB, E
Primary Accession	Q9NYJ7
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	64618
Antigen Region	519-548

Additional Information

Gene ID	10683
Other Names	Delta-like protein 3, Drosophila Delta homolog 3, Delta3, DLL3
Target/Specificity	This DLL3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 519-548 amino acids from the C-terminal region of human DLL3.
Dilution	IHC-P-Leica~~1:500 IF~~1:10~50 WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	DLL3 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	DLL3
Function	Inhibits primary neurogenesis. May be required to divert neurons along a specific differentiation pathway. Plays a role in the formation of somite boundaries during segmentation of the paraxial mesoderm (By similarity).
Cellular Location	Membrane; Single-pass type I membrane protein


Background

DLL3 encodes a member of the delta protein ligand family. This family functions as Notch ligands that are characterized by a DSL domain, EGF repeats, and a transmembrane domain.

References

Yerges,L.M. J. Bone Miner. Res. 24 (12), 2039-2049 (2009)
Heuss,S.F. Proc. Natl. Acad. Sci. U.S.A. 105 (32), 11212-11217 (2008)
Maisenbacher,M.K. Hum. Genet. 116 (5), 416-419 (2005)

Images



All lanes : Anti-DLL3 Antibody (C-term) at 1:2000 dilution
Lane 1: Human brain tissue lysate Lane 2: Mouse brain tissue lysate Lane 3: Rat brain tissue lysate
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 65 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.