10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# KAP1 (1A14) Rabbit Monoclonal Antibody

KAP1 (1A14) Rabbit Monoclonal Antibody
Catalog # AP93650

## Product Information

| | |
|---|---|
| **Application** | WB, IHC-P, IF, ICC |
| **Primary Accession** | Q13263 |
| **Reactivity** | Human, Mouse, Rat |
| **Clonality** | Monoclonal |
| **Calculated MW** | 88550 |

## Additional Information

| | |
|---|---|
| **Gene ID** | 10155 |
| **Other Names** | Transcription intermediary factor 1-beta, TIF1-beta, E3 SUMO-protein ligase TRIM28, 2.3.2.27, KRAB-associated protein 1, KAP-1, KRAB-interacting protein 1, KRIP-1, Nuclear corepressor KAP-1, RING finger protein 96, RING-type E3 ubiquitin transferase TIF1-beta, Tripartite motif-containing protein 28, TRIM28 (HGNC:16384), KAP1, RNF96, TIF1B |
| **Dilution** | WB~~1:1000 IHC-P~~N/A IF~~1:50~200 ICC~~N/A |
| **Storage Conditions** | -20℃ |

## Protein Information

| | |
|---|---|
| **Name** | TRIM28 ( HGNC:16384) |
| **Synonyms** | KAP1, RNF96, TIF1B |
| **Function** | E3 SUMO and ubiquitin ligase that plays a pivotal role in embryonic development, genomic imprinting, and maintenance of genomic stability through repression of repetitive and retroviral elements. Also involved in DNA repair, regulation of innate immunity or cellular energy homeostasis. Acts as a scaffold for assembling transcriptional repression complexes containing methyltransferases, histone deacetylases, and chromatin remodelers. Serves as a nuclear corepressor for KRAB domain-containing zinc finger proteins (KRAB-ZFPs), mediating gene silencing by recruiting CHD3, a subunit of the nucleosome remodeling and deacetylation (NuRD) complex, and SETDB1, which methylates histone H3 at 'Lys-9' (H3K9me), leading to heterochromatin formation. In collaboration with SETDB1, is also required for H3K9me3 and silencing of endogenous and introduced retroviruses in a DNA- methylation independent-pathway (By similarity). Functions as a coactivator for CEBPB and NR3C1 (glucocorticoid receptor) in transcriptional activation of ORM1, and as a corepressor for ERBB4. Inhibits E2F1 activity by promoting E2F1-HDAC1 complex formation and preventing E2F1 acetylation. This contributes to |

CDKN1A/p21(CIP1) regulation and provides a partial backup to suppress E2F1-mediated apoptosis in the absence of RB1. Mediates the nuclear localization of several KRAB-ZFP transcription factors including KOX1, ZNF268, and ZNF300 among others. In association with isoform 2 of ZFP90, is required for the transcriptional repressor activity of FOXP3 and the suppressive function of regulatory T-cells (Treg) (PubMed:23543754). Required to maintain a transcriptionally repressive state of genes in undifferentiated embryonic stem cells (ESCs) (PubMed:24623306). Acts as a corepressor for ZFP568. Beyond its nuclear functions, acts as a positive regulator of type I interferon (IFN-I) signaling by promoting Lys-63-linked ubiquitination of TBK1, facilitating TBK1-IRF3 complex formation (PubMed:38495890). Mediates TRAF6 SUMOylation to regulate its nucleo-cytoplasmic shuttling and modulate activation of the canonical NF-kappa-B signaling pathway (PubMed:39920527). Facilitates SUMOylation of NLRP3, protecting it from 'Lys-48'-linked ubiquitination and proteasomal degradation and thereby enhancing inflammasome activation (PubMed:34373456).


**Cellular Location**

Nucleus. Note=Associated with centromeric heterochromatin during cell differentiation through CBX1 (By similarity). Localizes to sites of DNA damage (PubMed:25593309) {ECO:0000250|UniProtKB:Q62318, ECO:0000269|PubMed:25593309}

**Tissue Location**

Expressed in all tissues tested including spleen, thymus, prostate, testis, ovary, small intestine, colon and peripheral blood leukocytes.

## Images




Western blot detection of KAP1 in K562,C6,3T3,Hela cell lysates using KAP1 antibody(1:1000 diluted).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.