10320 Camino Santa Fe, Suite G San Diego, CA 92121 Tel: 858.875.1900 Fax: 858.875.1999



# YAP1 (19W11) Rabbit Monoclonal Antibody

YAP1 (19W11) Rabbit Monoclonal Antibody Catalog # AP93664

### **Product Information**

**Application** WB, IHC, IF, ICC

Primary Accession P46937

Reactivity Human, Mouse
Clonality Monoclonal
Calculated MW 54462

## **Additional Information**

**Gene ID** 10413

Other Names Transcriptional coactivator YAP1, Yes-associated protein 1, Protein yorkie

homolog, Yes-associated protein YAP65 homolog, YAP1 (HGNC:16262), YAP65

**Dilution** WB~~1:1000 IHC~~1:100~500 IF~~1:50~200 ICC~~N/A

Storage Conditions -20°C

## **Protein Information**

Name YAP1 ( <u>HGNC:16262</u>)

Synonyms YAP65

**Function** Transcriptional regulator with dual roles as a coactivator and corepressor.

Critical downstream regulatory target in the Hippo signaling pathway, crucial for organ size control and tumor suppression by restricting proliferation and

promoting apoptosis (PubMed:<u>17974916</u>, PubMed:<u>18280240</u>,

PubMed: 18579750, PubMed: 21364637, PubMed: 30447097). The Hippo signaling pathway core involves a kinase cascade featuring STK3/MST2 and STK4/MST1, along with its regulatory partner SAV1, which phosphorylates and activates LATS1/2 in complex with their regulatory protein, MOB1. This activation leads to the phosphorylation and inactivation of the YAP1

oncoprotein and WWTR1/TAZ (PubMed:<u>18158288</u>). Phosphorylation of YAP1 by LATS1/2 prevents its nuclear translocation, thereby regulating the expression of its target genes (PubMed:<u>18158288</u>, PubMed:<u>26598551</u>, PubMed:<u>34404733</u>). The transcriptional regulation of gene expression

requires TEAD transcription factors and modulates cell growth,

anchorage-independent growth, and induction of epithelial- mesenchymal transition (EMT) (PubMed: 18579750). Plays a key role in tissue tension and 3D tissue shape by regulating the cortical actomyosin network, acting via ARHGAP18, a Rho GTPase activating protein that suppresses F-actin

polymerization (PubMed: 25778702). It also suppresses ciliogenesis by acting

as a transcriptional corepressor of TEAD4 target genes AURKA and PLK1 (PubMed: <u>25849865</u>). In conjunction with WWTR1, regulates TGFB1-dependent SMAD2 and SMAD3 nuclear accumulation (By similarity). Synergizes with WBP2 to enhance PGR activity (PubMed: <u>16772533</u>).


#### **Cellular Location**

Cytoplasm. Nucleus. Cell junction, tight junction {ECO:0000250|UniProtKB:A0A8C0NGY6}. Cell membrane. Note=Both phosphorylation and cell density can regulate its subcellular localization (PubMed:18158288, PubMed:20048001). Phosphorylation sequesters it in the cytoplasm by inhibiting its translocation into the nucleus (PubMed:18158288. PubMed:20048001, PubMed:34404733). At low density, predominantly nuclear and is translocated to the cytoplasm at high density (PubMed:18158288, PubMed:20048001, PubMed:25849865). PTPN14 induces translocation from the nucleus to the cytoplasm (PubMed:22525271). In the nucleus, phosphorylation by PRP4K induces nuclear exclusion (PubMed:29695716). Localized mainly to the nucleus in the early stages of embryo development with expression becoming evident in the cytoplasm at the blastocyst and epiblast stages (By similarity) Localizes to the cytoplasm and tight junctions following interaction with AMOT isoform 1 (PubMed:21205866). Localizes to tight junctions following interaction with AMOTL2 (By similarity). Translocates to the nucleus in the presence of SNAIL1 (By similarity). Found at the cell membrane in keratinocytes in response to mechanical strain (PubMed:31835537). {ECO:0000250|UniProtKB:A0A8C0NGY6, ECO:0000250|UniProtKB:P46938, ECO:0000269 | PubMed:18158288, ECO:0000269 | PubMed:20048001, ECO:0000269 | PubMed:21205866, ECO:0000269 | PubMed:22525271, ECO:0000269 | PubMed:25849865, ECO:0000269 | PubMed:29695716, ECO:0000269 | PubMed:31835537, ECO:0000269 | PubMed:34404733 }

#### **Tissue Location**

Increased expression seen in some liver and prostate cancers. Isoforms lacking the transactivation domain found in striatal neurons of patients with Huntington disease (at protein level).

# **Images**



Western blot detection of YAP1 in Hela cell lysates using YAP1 antibody(1:1000 diluted).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.