

Phospho-Smad1/5/9 (S463/S465/S467) (11L16) Rabbit Monoclonal Antibody

Phospho-Smad1/5/9 (S463/S465/S467) (11L16) Rabbit Monoclonal Antibody
Catalog # AP93680

Product Information

| | |
|-------------------|------------------------|
| Application | WB |
| Primary Accession | Q99717 |
| Reactivity | Rat, Human, Mouse |
| Clonality | Monoclonal |
| Calculated MW | 52258 |

Additional Information

| | |
|--------------------|---|
| Gene ID | 4090 |
| Other Names | Mothers against decapentaplegic homolog 5, MAD homolog 5, Mothers against DPP homolog 5, JV5-1, SMAD family member 5, SMAD 5, Smad5, hSmad5, SMAD5, MADH5 |
| Dilution | WB~~1:1000 |
| Storage Conditions | -20°C |

Protein Information


| | |
|-------------------|---|
| Name | SMAD5 |
| Synonyms | MADH5 |
| Function | <p>Transcriptional regulator that plays a role in various cellular processes including embryonic development, cell differentiation, angiogenesis and tissue homeostasis (PubMed:12064918, PubMed:16516194). Upon BMP ligand binding to their receptors at the cell surface, is phosphorylated by activated type I BMP receptors (BMPRI) and associates with SMAD4 to form a heteromeric complex which translocates into the nucleus acting as transcription factor (PubMed:9442019). In turn, the hetero-trimeric complex recognizes cis- regulatory elements containing Smad Binding Elements (SBEs) to modulate the outcome of the signaling network (PubMed:33510867). Non-phosphorylated SMAD5 has a cytoplasmic role in energy metabolism regulation by promoting mitochondrial respiration and glycolysis in response to cytoplasmic pH changes (PubMed:28675158). Mechanistically, interacts with hexokinase 1/HK1 and thereby accelerates glycolysis (PubMed:28675158).</p> |
| Cellular Location | Cytoplasm. Nucleus Mitochondrion. Note=Cytoplasmic in the absence of |

ligand. Migrates to the nucleus when complexed with SMAD4

Tissue Location

Ubiquitous.

Images



Western blot analysis of Phospho-Smad5 (S463/S465) expression in (1) HeLa cell lysate; (2) HeLa cell treated with BMP-4 lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.