

EIF2S1 (18K9) Rabbit Monoclonal Antibody

EIF2S1 (18K9) Rabbit Monoclonal Antibody

Catalog # AP93737

Product Information

Application	WB, IHC-P, IP, IF
Primary Accession	P05198 , Q6ZWX6 , P68101
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Calculated MW	36112

Additional Information

Gene ID	1965
Dilution	WB~~1:1000 IHC-P~~N/A IP~~N/A IF~~1:50~200
Storage Conditions	-20°C

Protein Information

Name	EIF2S1 (HGNC:3265)
Synonyms	EIF2A
Function	Member of the eIF2 complex that functions in the early steps of protein synthesis by forming a ternary complex with GTP and initiator tRNA (PubMed:16289705, PubMed:38340717). This complex binds to a 40S ribosomal subunit, followed by mRNA binding to form a 43S pre- initiation complex (43S PIC) (PubMed:16289705). Junction of the 60S ribosomal subunit to form the 80S initiation complex is preceded by hydrolysis of the GTP bound to eIF2 and release of an eIF2-GDP binary complex (PubMed:16289705). In order for eIF2 to recycle and catalyze another round of initiation, the GDP bound to eIF2 must exchange with GTP by way of a reaction catalyzed by eIF2B (PubMed:16289705). EIF2S1/eIF2-alpha is a key component of the integrated stress response (ISR), required for adaptation to various stress: phosphorylation by metabolic-stress sensing protein kinases (EIF2AK1/HRI, EIF2AK2/PKR, EIF2AK3/PERK and EIF2AK4/GCN2) in response to stress converts EIF2S1/eIF2-alpha in a global protein synthesis inhibitor, leading to an attenuation of cap-dependent translation, while concomitantly initiating the preferential translation of ISR-specific mRNAs, such as the transcriptional activators ATF4 and QRI1, and hence allowing ATF4- and QRI1-mediated reprogramming (PubMed:19131336, PubMed:33384352, PubMed:38340717). EIF2S1/eIF2-alpha also acts as an activator of mitophagy in response to mitochondrial damage: phosphorylation by EIF2AK1/HRI promotes relocalization to the mitochondrial surface, thereby triggering PRKN-independent mitophagy (PubMed:38340717).


Cellular Location

Cytoplasm, Stress granule {ECO:0000250|UniProtKB:Q6ZWX6}. Cytoplasm, cytosol {ECO:0000250|UniProtKB:P56286}. Mitochondrion. Note=Colocalizes with NANOS3 in the stress granules (By similarity). Relocalizes to the surface of mitochondria in response to mitochondrial damage and phosphorylation by EIF2AK1/HRI (PubMed:38340717). {ECO:0000250|UniProtKB:Q6ZWX6, ECO:0000269|PubMed:38340717}

Background

The translation initiation factor EIF2 catalyzes the first regulated step of protein synthesis initiation, promoting the binding of the initiator tRNA to 40S ribosomal subunits. Binding occurs as a ternary complex of methionyl-tRNA, EIF2, and GTP. EIF2 is composed of 3 nonidentical subunits, the 36-kD EIF2-alpha subunit (EIF2S1), the 38-kD EIF2-beta subunit (EIF2S2; MIM 603908), and the 52-kD EIF2-gamma subunit (EIF2S3; MIM 300161). The rate of formation of the ternary complex is modulated by the phosphorylation state of EIF2-alpha (Ernst et al., 1987 [PubMed 2948954]).[supplied by OMIM, Feb 2010]

Images



Western blot analysis of extracts from MCF-7, HepG2 cells using AP93737 at 1:1000.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.